

# Contents:

## The Journal of Cell Biology

Volume 107, Number 2, August 1988

- 397 **The fates of chicken nuclear lamin proteins during mitosis: evidence for a reversible redistribution of lamin B<sub>2</sub> between inner nuclear membrane and elements of the endoplasmic reticulum.** R. Stick, B. Angres, C. F. Lehner, and E. A. Nigg
- 407 **Localization of protamine 1 mRNA in different stages of the cycle of the rat seminiferous epithelium.** P. Mali, M. Sandberg, E. Vuorio, P. C. Yelick, N. B. Hecht, and M. Parvinen
- 413 **Unequal synthesis and differential degradation of  $\alpha$  and  $\beta$  spectrin during murine erythroid differentiation.** M. E. Lehnert and H. F. Lodish
- 427 **The use of monospecific antisera and cRNA probes to localize the major changes in keratin expression during normal and abnormal epidermal differentiation.** A. Stoler, R. Kopan, M. Duvic, and E. Fuchs
- 447 **Human interleukin 1 $\beta$  is not secreted from hamster fibroblasts when expressed constitutively from a transfected cDNA.** P. R. Young, D. J. Hazuda, and P. L. Simon
- 457 **Signals for the incorporation and orientation of cytochrome P450 in the endoplasmic reticulum membrane.** S. Monier, P. Van Luc, G. Kreibich, D. D. Sabatini, and M. Adesnik
- 471 **A specific sorting signal is not required for the polarized secretion of newly synthesized proteins from cultured intestinal epithelial cells.** M. J. Rindler and M. G. Traber
- 481 **Coupling membranes as energy-transmitting cables. I. Filamentous mitochondria in fibroblasts and mitochondrial clusters in cardiomyocytes.** A. A. Amchenkova, L. E. Bakeeva, Y. S. Chentsov, V. P. Skulachev, and D. B. Zorov
- 497 **Coupling membranes as energy-transmitting cables. II. Cyanobacterial trichomes.** I. I. Severina, V. P. Skulachev, and D. B. Zorov
- 503 **Topogenesis of mitochondrial inner membrane uncoupling protein. Rerouting transmembrane segments to the soluble matrix compartment.** X. Liu, A. W. Bell, K. B. Freeman, and G. C. Shore
- 511 **Identification of diphtheria toxin receptor and a nonproteinous diphtheria toxin-binding molecule in *Vero* cell membrane.** E. Mekada, Y. Okada, and T. Uchida
- 521 **Signal transduction in *Dictyostelium* FGD A mutants with a defective interaction between surface cAMP receptor and a GTP-binding regulatory protein.** F. Kesbeke, B. E. Snaar-Jagalska, and P. J. M. Van Haastert
- 529 **Concurrent changes in *Dunaliella salina* ultrastructure and membrane phospholipid metabolism after hyperosmotic shock.** K. J. Einspahr, M. Maeda, and G. A. Thompson, Jr.
- 539 **Evidence for involvement of metalloendoproteases in a step in sea urchin gamete fusion.** J. L. Roe, H. A. Farach, Jr., W. J. Strittmatter, and W. J. Lennarz
- 545 **Immunolocalization of meta-vinculin in human smooth and cardiac muscles.** A. M. Belkin, O. I. Ornatsky, M. A. Glukhova, and V. E. Koteliansky
- 555 **Localization of phospholamban in smooth muscle using immunogold electron microscopy.** D. G. Ferguson, E. F. Young, L. Raeymaekers, and E. G. Kranias
- 563 **Differential localization of tropomyosin isoforms in cultured nonmuscle cells.** J. J.-C. Lin, T. E. Hegmann, and J. L.-C. Lin
- 573 **Analysis of the upstream regions governing expression of the chicken cardiac troponin T gene in embryonic cardiac and skeletal muscle cells.** J. H. Mar, P. B. Antin, T. A. Cooper, and C. P. Ordahl
- 587 **Posttranslational incorporation of contractile proteins into myofibrils in a cell-free system.** M. Bouché, S. M. Goldfine, and D. A. Fischman
- 597 **Three-dimensional reconstruction of an actin bundle.** E. S. A. Bullitt, D. J. DeRosier, L. M. Coluccio, and L. G. Tilney

Contents continued on reverse of this cover

- 613 Two classes of actin microfilaments are associated with the inner cytoskeleton of axons. K. R. Fath and R. J. Lasek
- 623 The mechanism of anaphase spindle elongation: uncoupling of tubulin incorporation and microtubules sliding during in vitro spindle reactivation. H. Masuda, K. L. McDonald, and W. Z. Cande
- 635 The centrin-based cytoskeleton of *Chlamydomonas reinhardtii*: distribution in interphase and mitotic cells. J. L. Salisbury, A. T. Baron, and M. A. Sanders
- 643 PC12 cells express juvenile microtubule-associated proteins during nerve growth factor-induced neurite outgrowth. B. Brugg and A. Matus
- 651 Microtubule dynamics in nerve cells: analysis using microinjection of biotinylated tubulin into PC12 cells. S. Okabe and N. Hirokawa
- 665 Tension and compression in the cytoskeleton of PC-12 neurites II: quantitative measurements. T. J. Dennerll, H. C. Joshi, V. L. Steel, R. E. Buxbaum, and S. R. Heidemann
- 675 Distribution of the myelin-associated glycoprotein and P<sub>0</sub> protein during myelin compaction in quaking mouse peripheral nerve. B. D. Trapp
- 687 Isolation and characterization of a laminin-binding protein from rat and chick muscle. D. E. Hall, K. A. Frazer, B. C. Hann, and L. F. Reichardt
- 699 Amino acid sequence and distribution of mRNA encoding a major skeletal muscle laminin binding protein: an extracellular matrix-associated protein with an unusual COOH-terminal polyaspartate domain. D. O. Clegg, J. C. Helder, B. C. Hann, D. E. Hall, and L. F. Reichardt
- 707 Distribution and role in regeneration of N-CAM in the basal laminae of muscle and Schwann cells. F. Rieger, M. Nicolet, M. Pincon-Raymond, M. Murawsky, G. Levi, and G. M. Edelman
- 721 Type VIII collagen has a restricted distribution in specialized extracellular matrices. R. Kapoor, L. Y. Sakai, S. Funk, E. Roux, P. Bornstein, and E. H. Sage
- 731 The extracellular matrix of *Xenopus laevis* eggs: a quick-freeze, deep-etch analysis of its modification at fertilization. C. A. Larabell and D. E. Chandler
- 743 Endothelial cell-derived heparan sulfate binds basic fibroblast growth factor and protects it from proteolytic degradation. O. Saksela, D. Moscatelli, A. Sommer, and D. B. Rifkin
- 753 Metabolism of receptor-bound and matrix-bound basic fibroblast growth factor by bovine capillary endothelial cells. D. Moscatelli
- 761 Cell surface fibroblast growth factor and epidermal growth factor receptors are permanently lost during skeletal muscle terminal differentiation in culture. B. B. Olwin and S. D. Hauschka
- 771 Transforming growth factor- $\beta$ -induced growth inhibition and cellular hypertrophy in cultured vascular smooth muscle cells. G. K. Owens, A. A. T. Geisterfer, Y. W.-H. Yang, and A. Komoriya
- 781 Tumor necrosis factor and immune interferon act in concert to slow the lateral diffusion of proteins and lipids in human endothelial cell membranes. A. H. Stolpen, D. E. Golan, and J. S. Pober
- 791 Two independent mechanisms for escaping epidermal growth factor-mediated growth inhibition in epidermal growth factor receptor-hyperproducing human tumor cells. M. Hirai, S. Gamou, S. Minoshima, and N. Shimizu
- 801 Anomalous binding of epidermal growth factor to A431 cells is due to the effect of high receptor densities and a saturable endocytic system. H. S. Wiley
- 811 Focal expression of insulin-like growth factor I in rat kidney collecting duct. J. D. Bortz, P. Rotwein, D. DeVol, P. J. Bechtel, V. A. Hansen, and M. R. Hammerman
- 821 Correction

---

Cover picture: Light micrograph, Nomarski optics, of a *Dunaliella salina* cell growing without osmotic stress in 1.7 M NaCl-containing medium. Micrograph courtesy of Randy Wayne and John LaClaire. See related article in this issue by Einspahr et al., pp. 529-538.