

Prospore Membrane Formation Defines a Developmentally Regulated Branch of the Secretory Pathway in Yeast

Aaron M. Neiman

Department of Biochemistry and Cell Biology, State University of New York, Stony Brook, Stony Brook, New York 11794

Abstract. Spore formation in yeast is an unusual form of cell division in which the daughter cells are formed within the mother cell cytoplasm. This division requires the de novo synthesis of a membrane compartment, termed the prospore membrane, which engulfs the daughter nuclei. The effect of mutations in late-acting genes on sporulation was investigated. Mutation of *SEC1*, *SEC4*, or *SEC8* blocked spore formation, and electron microscopic analysis of the *sec4-8* mutant indicated that this inability to produce spores was caused by a failure to form the prospore membrane. The soluble NSF attachment protein 25 (SNAP-25) homologue *SEC9*, by contrast, was not required for sporulation.

The absence of a requirement for *SEC9* was shown to be due to the sporulation-specific induction of a second, previously undescribed, SNAP-25 homologue, termed *SPO20*. These results define a developmentally regulated branch of the secretory pathway and suggest that spore morphogenesis in yeast proceeds by the targeting and fusion of secretory vesicles to form new plasma membranes in the interior of the mother cell. Consistent with this model, the extracellular proteins Gas1p and Cts1p were localized to an internal compartment in sporulating cells. Spore formation in yeast may be a useful model for understanding secretion-driven cell division events in a variety of plant and animal systems.

SPORULATION in *Saccharomyces cerevisiae* is an unusual form of cell division in which the daughter cells are formed within the original mother cell (Esposito and Klapholz, 1981). In this process, a single round of DNA replication followed by two meiotic divisions occurs within a single, intact nuclear envelope. During the second meiotic division, an extension of the outer plaque is formed on the cytoplasmic side of each of the four spindle pole bodies (Moens, 1971; Moens and Rapport, 1971; Byers, 1981). A flattened membrane sac then forms on the cytoplasmic face of each of the outer plaques. This membrane, which is sometimes called the prospore wall, is referred to in this work as the prospore membrane. As meiosis II progresses, the chromatin segregates into four lobes of the nucleus and each prospore membrane grows larger to engulf the adjacent nuclear lobe. As meiosis II is completed and the nucleus divides, each prospore membrane fuses with itself so that every daughter nucleus (and associated cytoplasm) is captured inside two, now distinct, unit membranes of the prospore membrane, creating immature spores. The bilayer of the prospore membrane closest to the daughter nucleus now serves as the plasma membrane of the spore. In the final step of spore morphogenesis, the spore wall is formed in the luminal space between the two

membranes derived from the prospore membrane (Lynn and Magee, 1970).

Although the cytology of spore morphogenesis has been described, little is known about the molecular and genetic requirements for spore formation. A growing number of mutants have been identified that proceed through meiosis II but do not form proper spores (Moens et al., 1974; Briza et al., 1990; Friesen et al., 1994; Krisak et al., 1994; Rose et al., 1995; Tu et al., 1996). Some of the mutations disrupt proper assembly of the spore wall (Briza et al., 1990; Friesen et al., 1994; Krisak et al., 1994), whereas others seem to disrupt the coordination between prospore membrane growth and nuclear division. A *spo3* mutant, for example, suffers from a delay in meiosis II, the consequence of which is premature closure (relative to nuclear division) of the prospore membrane and, therefore, the failure to package the daughter nuclei into spores (Moens et al., 1974).

It has been suggested on morphological grounds that the prospore membrane is a derivative of the ER (Lynn and Magee, 1970). However, this proposition has not been directly tested. One means to investigate the origin of the prospore membrane is to examine the effect of mutations in *SEC* genes on development of this membrane. A large number of *sec* mutants that block the transit of vesicles through the secretory pathway at distinct steps have been isolated in *S. cerevisiae* (Novick et al., 1981). One subset of *SEC* genes, for instance, are specifically required at the end of the secretory pathway for the fusion of post-Golgi

Address all correspondence to Aaron M. Neiman, Department of Biochemistry and Cell Biology, State University of New York, Stony Brook, Stony Brook, NY 11794. Tel.: (516) 632-8565. Fax: (516) 632-8575. E-mail: aneiman@mcbgsi.bio.sunysb.edu

vesicles with the plasma membrane (Novick et al., 1981). These late-acting *SEC* genes include *SEC1*, 2, 3, 4, 5, 6, 8, 9, 10, and 15, as well as two pairs of redundant genes *SSO1* and 2 and *SNC1* and 2. The molecular function of many of these gene products is understood. *SEC4*, for example, encodes a GTPase of the rab family (Salminen and Novick, 1987). *SEC9* encodes a subunit of the plasma membrane t-SNARE complex (Aalto et al., 1993; Brennwald et al., 1994). Additionally, Sec6p, Sec8p, and Sec15p are subunits of a large, cytoplasmic protein complex that localizes to sites of vesicle fusion on the plasma membrane, and several of the other late-acting *SEC* genes are required for proper assembly of this complex (Terbush and Novick, 1995).

A large body of evidence supports a general model for vesicle fusion events in the secretory pathway, termed the soluble NSF attachment protein receptor (SNARE)¹ hypothesis, in which fusion of transport vesicles with the appropriate target membrane is controlled by SNARE complexes (Söllner et al., 1993; Sjøgaard et al., 1994; Rothman, 1994). According to this model, every vesicle carries on its surface a vesicular (v-)SNARE that can only interact with target (t-)SNARE found on the correct acceptor membrane. Interaction of SNAREs is further regulated by the action of rab-family GTPases. The specificity of SNARE interactions controls the specificity of vesicular transport in the cell. In the case of secretion in *S. cerevisiae*, vesicles destined for the plasma membrane carry on their surface the v-SNARE proteins Snc1p or Snc2p (Protopopov et al., 1993). These Snc proteins are predicted to interact specifically with the t-SNARE proteins Sec9p and Sso1p or Sso2p, ensuring that Snc-containing vesicles fuse only with the plasma membrane and that no other vesicles fuse with the plasma membrane. The association of these proteins is expected to be regulated by Sec4p. Indeed, *SEC4* has genetic interactions with *SEC9* and, under appropriate conditions, Sso1p, Snc1p, and Sec9p can be coimmunoprecipitated (Brennwald et al., 1994).

Because the prospore membrane functions as the plasma membrane of the spore, it seemed possible that, like new plasma membrane, the prospore membrane was derived from post-Golgi vesicles, rather than directly from the ER as suggested by Lynn and Magee (1970). To differentiate between these possibilities, mutations in late-acting *SEC* genes, which effect only post-Golgi vesicular transport, were examined for their effect on spore formation. The results identify a previously unknown branch of the secretory pathway in yeast and suggest that spore morphogenesis begins by the de novo synthesis of a new plasma membrane in the interior of the mother cell.

Materials and Methods

Yeast Strains and Media

S. cerevisiae strains used in this work are listed in Table I. To make strains AN79, AN81, AN126, and AN127, haploids AN63-2C, AN61-1A, HSF187, and AN65-1C (*sec9-4*, *sec4-8*, *sec8-9*, and *sec1-1*, respectively) were crossed to the SK1-related strain S2683 to create diploids AN74,

AN75, AN124, and AN125. Several *sec* segregants from these diploids were crossed to each other to find diploids that sporulate efficiently at permissive temperature (AN79 AN81, AN126, and AN127). To create the *spo20ΔURA3* diploid AN67, *SPO20* was first disrupted in S2683 and RKY1145 by one-step gene replacement (Rothstein, 1983) using EcoRI- and XbaI-digested *pspo20ΔURA3*. The resulting haploids (AN1037 and AN1038) were then mated to produce AN67. AN80 was constructed in an identical manner except that the haploid strains used were AN74-3A and AN74-7C. Unless otherwise noted, standard media and growth conditions were used (Rose and Fink, 1990).

Sporulation Assays

Strains were sporulated by overnight growth in YPD followed by 1:300 dilution into YPacetate medium. Cultures were then grown overnight to midlog phase, pelleted, washed once, and resuspended in 2% KOAc medium at a concentration of 3×10^7 cells/ml. For strains harboring temperature-sensitive mutations, cultures were incubated in sporulation medium at 23°C for ~2 h before shift to 34°C. At intervals, aliquots were removed and fixed by addition of formaldehyde to a final concentration of 3.7%. Samples were then stained with 4',6'-diamidino-2-phenylindolehydrochloride (DAPI) and observed using a Zeiss Axiophot microscope (Thornwood, NY).

Plasmids

The *SPO20* region was cloned directly from genomic DNA by PCR using the primers ANO85 (5'-GGC TGA ATT CTA GGC GCT TTC AAC C) and ANO86 (5'-GCC GTC TAG AGT GTA TAA CAG ATC ACC) and *Taq* polymerase (Boehringer Mannheim Corp., Indianapolis, IN). The resulting fragment was digested with EcoRI and XbaI and cloned into EcoRI-XbaI-cut Bluescript SK⁻ (Promega Corp., Madison, WI) to create pSK-SEC9H. The *spo20ΔURA3* allele was created by replacing a BamHI-BglII fragment of pSK-SEC9H carrying the central 0.7 kb of the *SPO20* coding region with a 1.1-kb BamHI-BglII fragment of pMH3 (a gift of N. Hollingsworth, SUNY Stony Brook) carrying the *URA3* gene. The resulting plasmid *pspo20ΔURA3* was digested with EcoRI and XbaI before transformation to release the disruption fragment. The *SPO20-HA* fusion allele was constructed in three steps. First, the EcoRI-XbaI fragment of pSK-SEC9H carrying the entire *SPO20* region was cloned into pRS315 (Sikorski and Hieter, 1989). Using the primers ANO95 (5'-GTG CTG AAT TCT ATA TAA TGG GGT TCA G) and ANO96 (5'-GTG TAT AAC TCA TAT GCA TCT TTT CCC G) and pSK-SEC9H as a template, PCR was then used to introduce EcoRI and NsiI sites 5' of the ATG and at the stop codon of the *SPO20* open reading frame, respectively. This PCR fragment was digested with EcoRI and NsiI and cloned into EcoRI-PstI-cut pSKHA3 (Neiman et al., 1997) to create an in-frame fusion of three copies of the influenza hemagglutinin epitope (YPYDVPDY) to the carboxy terminus of *SPO20*. Finally, a BglII-XbaI fragment coding for the COOH-terminal portion of the Spo20-HA fusion protein was swapped with a corresponding BglII-XbaI fragment of pRS315-SPO20 to create pRS315-SPO20-3xHA. A NotI-XhoI fragment of pRS315-SPO20-3xHA was cloned into similarly cut pRS425 to create pRS425-SPO20-3xHA. The original PCR cloning from the genome did not introduce deleterious mutations into the *SPO20* coding region, as shown by the fact that pRS315-SPO20 completely complements the *spo20Δ* sporulation defect of strain AN67. The hemagglutinin (HA) tag may mildly impair Spo20p function, as a *spo20ΔURA3* strain carrying pRS315-SPO20-3xHA is reduced in sporulation ~50% relative to the same strain carrying pRS315-SPO20.

To examine whether *SEC9* can rescue the *spo20* sporulation defect, a genomic clone of *SEC9* was isolated by complementation of the *sec9-4* temperature-sensitive phenotype with a YEpl3 based library and then transformed into strain AN67. For the reciprocal experiment, the PCR product carrying the *SPO20* coding region (generated using primers AN95 and ANO96, see above) was cloned as a blunt end fragment into EcoRV-cut Bluescript KS⁺ and then subcloned as a 1.2-kb EcoRI-SalI fragment into plasmid pRD53 (R. Deshaies), placing *SPO20* under the control of the *GAL1* promoter. This plasmid, pRD53-SPO20, was then transformed into strain AN63-2C, and growth was assessed on galactose plates incubated at 36°C.

Electron Microscopy

Samples were prepared for the electron microscope by a modification of the procedure of Moens (1971). Samples were fixed by addition of glu-

1. Abbreviations used in this paper: DAPI, 4',6'-diamidino-2-phenylindolehydrochloride; HA, hemagglutinin; NSF, N-ethylmaleimide-sensitive factor; SNAP, soluble NSF attachment protein; SNARE, SNAP receptor; v- and t-SNARE, vesicular and target SNARE.

Table I. Strains

Strain	Genotype	Source
AN61-1A	<i>MATa ura3-52 sec4-8</i>	this work [‡]
AN63-2C	<i>MATa ura3-52 sec9-4</i>	this work [§]
HSF187	<i>MATa sec8-9</i>	N. Dean
AN65-1C	<i>MATα ura3 sec1-1</i>	this work
RKY1145	<i>MATa ura3 leu2ΔhisG his4-x lys2 hoΔLYS2</i>	N. Hollingsworth
S2683	<i>MATα ura3 leu2-k arg4-NspI lys2 hoΔLYS2</i>	N. Hollingsworth
AN1037	as RKY1145, plus <i>spo20ΔURA3</i>	see text
AN1038	as S2683, plus <i>spo20ΔURA3</i>	see text
AN74-3A*	<i>MATa ura3 arg4-NspI sec9-4</i>	segregant of AN63-2C × S2683
AN74-7C*	<i>MATα ura3 leu2-k sec9-4</i>	segregant of AN63-2C × S2683
NH144	<i>MATa/MATα ura3/ura3/ura3 leu2ΔhisG/leu2-k his4-x/+ his4-x/+ arg4-NspI/+ lys2/lys2 hoΔLYS2/hoΔLYS2</i>	RKY1145 × S2683
AN67	as NH144 plus <i>spo20ΔURA3/spo20ΔURA3</i>	AN1037 × AN1038
AN79*	<i>MATa/MATα ura3/ura3 leu2-k/+ arg4-NspI/+ sec9-4/sec9-4</i>	AN74-3A × AN74-7C
AN80*	as AN79, plus <i>spo20ΔURA3/spo20ΔURA3</i>	see text
AN81*	<i>MATa/MATα ura3/ura3 leu2-k/+ arg4-NspI/+ lys2/+ sec4-8/sec4-8</i>	see text
AN126*	<i>MATa/MATα ura3/ura3 leu2-k/+ arg4-NspI/+ sec8-9/sec8-9</i>	see text
AN127*	<i>MATa/MATα ura3/ura3 leu2-k/+ arg4-NspI/+ sec1-1/sec1-1</i>	see text

*These strains are Lys⁺, which may be due to any of the following genotypes: *LYS2 ho*, *lys2 ho::LYS2*, or *LYS2 ho::LYS2*.

[‡]Segregant from an outcross of strain NY774 (*sec4-8*) from D. Terbush.

[§]Segregant from an outcross of strain CKY31 (*sec9-4*) from C. Kaiser.

^{||}Segregant from an outcross of strain CKY23 (*sec1-1*) from C. Kaiser.

taraldehyde (Sigma Chemical Co., St. Louis, MO) directly to the culture medium to a final concentration of 2.5%, mixed, and incubated on ice overnight. They were then washed twice in distilled water before being resuspended in 4% KMnO₄ for 30 min. The KMnO₄ was then removed by washing with distilled water and the samples were resuspended in a saturated uranyl acetate solution for 30 min. After one wash with distilled water, the samples were dehydrated by a series of ethanol washes and a final wash in 100% propylene oxide. The samples were then washed three times in Epon mix (5 ml Epon-812, 1.5 ml dodecyl succinic anhydride [DDSA], 3.5 ml nadic methyl anhydride [NMA]). Finally, the samples were resuspended in Epon mix with DMP-30 (0.15 ml), left for 2 d at room temperature, and then shifted to 60°C for 24 h before sectioning. For all strains described here, at least 50 cells were analyzed that were, as judged by nuclear morphology, postmeiotic or in meiosis II.

Invertase and Carboxypeptidase Y Assays

Invertase activity was monitored as described (Pelham et al., 1988). Carboxypeptidase Y sorting and processing was monitored by Western analysis and colony lifts using anti-CPY antibodies (N. Dean, SUNY Stony Brook) as described previously (Rothman et al., 1986).

Western Analysis

Strain AN67 carrying pRS425-SPO20-3×HA was sporulated in liquid medium, and at intervals aliquots were removed. Extracts were made by disruption with glass beads in lysis buffer (10 mM Hepes, pH 7.5, 200 mM KCl, 0.1% NP-40) followed by 30 min of centrifugation in a microfuge at 13,000 rpm. Samples were separated by SDS-PAGE in a 10% gel and then blotted to nitrocellulose. Filters were probed with the anti-HA antibody 12CA5 (BAbCo, Berkeley, CA) at a 1:3,000 dilution and then HRP-conjugated anti-mouse secondary antibodies. Bands were visualized using ECL (Amersham Corp., Arlington Heights, IL).

Indirect Immunofluorescence

Immunofluorescence was performed essentially as described (Neiman et al., 1997). The anti-Gas1p and anti-Cts1p antibodies (N. Dean) were used at a 1:1 dilution. The secondary antibodies used were goat anti-rabbit coupled to Cy3 (Cappel Laboratories, Malvern, PA).

Results

Late-acting SEC Genes Are Required for Spore Formation

To determine if late-acting *SEC* genes play a role in spore morphogenesis, diploids homozygous for the temperature-sensitive mutations *sec1-1*, *sec4-8*, *sec8-9*, or *sec9-4* were tested for their ability to form spores. Strains carrying these mutations were incubated briefly in sporulation medium at permissive temperature (23°C) and then shifted to nonpermissive temperature (34°C). Progression through meiosis was scored by staining the cells with DAPI and monitoring the appearance of tetranucleate staining by fluorescence microscopy, while formation of spores was assayed by phase microscopy. All four strains sporulated efficiently at permissive temperature (Table II). However, strains homozygous for *sec1-1*, *sec4-8*, or *sec8-9* were severely impaired in sporulation at the restrictive temperature. DAPI staining indicated that the *sec1-1/sec1-1*, *sec4-8/sec4-8*, and *sec8-9/sec8-9* strains formed tetranucleate cells nearly as well as wild type (NH144) (Table II). Some tetranucleate staining may represent segregation of the chromatin within a single nucleus rather than the completion of nuclear division (Kupiec et al., 1997). Nonetheless, this result suggests that the failure of these mutants to sporulate is an inability to form spores, per se, rather than a defect in the completion of meiosis.

In contrast, the *sec9-4/sec9-4* strain (AN79) sporulated well at the restrictive temperature (Table II). This difference between *sec9-4* and the other *sec* mutants is apparently not due to leakiness of the *sec9-4* allele since in all *sec* strains tested the nonpermissive temperature used for

Table II. Sporulation in *sec* Mutant Strains

Strain	Relevant genotype	Percent asci 23°C*	Percent asci 34°C*	Percent tetranucleate 34°C‡
NH144	wild type	96	95	61
AN127	<i>sec1-1/sec1-1</i>	80	<0.2	35
AN81	<i>sec4-8/sec4-8</i>	74	0.4	46
AN126	<i>sec8-9/sec8-9</i>	80	<0.2	39
AN79	<i>sec9-4/sec9-4</i>	63	69	23
AN67	<i>spo20Δ/spo20Δ</i>	<0.2	<0.2	32

*Cultures were transferred to sporulation medium, incubated for 3 h at 23°C, and then split in two. One half was left at 23°C, and the other was shifted to 34°C. After overnight incubation, all cultures were examined by phase contrast microscopy. For each culture, at least 500 cells were counted.

‡Tetranucleate staining was unstable in strains AN67, AN81, AN126, and AN127, probably because of the failure to package nuclei (see text). Therefore cells were fixed 4 h after shift to 34°C.

sporulation was above the vegetative restrictive temperature (Salminen and Novick, 1987; Neiman, A.M., unpublished observations).

sec4/sec4 Diploids Do Not Form Prospore Membranes

The failure of *sec1-1*, *sec4-8*, and *sec8-9* mutants to form visible spores could be due to a defect at any point in the development of refractile spores. For instance, these mutant strains might form the prospore membrane normally but be unable to form a mature spore wall. To examine where in the process of spore morphogenesis the *sec4* mutant defect occurs, the *sec4-8* homozygous diploid AN81 was compared with a *SEC4* diploid by transmission electron microscopy. Cultures of AN81 and NH144 were incubated in sporulation medium at 23°C for 3 h and then shifted to 34°C. At intervals, cells were removed and prepared for electron microscopy as described in Materials and Methods. Fig. 1 *A* shows a typical wild-type cell at a late stage of prospore membrane growth. The prospore membrane is a semicircular structure centered near the spindle pole body and curving down around a lobe of the nucleus. At a later stage (Fig. 1 *B*), the prospore membrane has closed around a daughter nucleus and a spore wall is forming. By contrast, in the *sec4-8/sec4-8* diploid, membranes resembling the prospore membrane were never observed. In cells that appeared to be in meiosis II, as judged by their nuclear morphology (Fig. 1 *C*), no semicircular membranes were evident, and at later stages, cells appeared to have multiple, unpackaged daughter nuclei (Fig. 1 *D*). These results indicate that in the *sec4-8* mutant, the prospore membrane does not form. Given the known function of *SEC4* in the fusion of post-Golgi secretory vesicles with the plasma membrane, these results suggest that the prospore membrane is formed by the fusion of secretory vesicles within the cytoplasm of the cell.

A Second Soluble NSF Attachment Protein (SNAP-25) Homologue Functions in Spore Morphogenesis

A number of genetic interactions between *SEC4* and *SEC9* have led to the suggestion that Sec9p is the effector for Sec4p in secretory vesicle fusion with the plasma membrane (Brennwald et al., 1994). Therefore, the observation that *SEC4* but not *SEC9* is required for prospore membrane formation might suggest that fusion of vesicles with the prospore membrane is mechanistically different from




Figure 1. Prospore membranes are absent in a *sec4/sec4* mutant. (A) *SEC4* cell (NH144) late in prospore membrane formation. The gap in the nuclear envelope underneath the outer plaque is the position of the spindle pole body. (B) NH144, after closure. (C and D) Cells of strain AN81 (*sec4-8/sec4-8*) at stages comparable to those in A and B, respectively. N, nucleus; M, mitochondrion; V, vacuole; OP, outer plaque; PrM, prospore membrane; SW, spore wall. All cells were sporulated at 23°C for 3 h and then shifted to the restrictive temperature for *sec4-8* (34°C). Bars: (A and C) 200 nm; (B and D) 500 nm.

fusion with the plasma membrane. Alternatively, there may be some sporulation-specific protein that can replace Sec9p in prospore membrane formation. To address this second possibility, the yeast genome was screened for *SEC9*-related sequences.

The *S. cerevisiae* genome contains one open reading frame, YMR017w, with significant sequence similarity to *SEC9*. For reasons described below, this open reading frame has been designated *SPO20*. The predicted protein encoded by *SPO20* is 397 amino acids in length with 37% identity and 55% similarity to Sec9p over the carboxy-terminal 223 residues (Fig. 2). The amino-terminal region of Spo20p has no similarity to Sec9p or any other sequence in GenBank. In the domain conserved between Spo20p and Sec9p, Spo20p contains two 60 amino acid regions that together show 28% identity and 50% similarity to the mammalian t-SNARE subunit SNAP-25. Thus, *SPO20* encodes a second *S. cerevisiae* SNAP-25 homologue.

The *SPO20* gene was cloned by PCR from yeast genomic DNA, and a diploid strain homozygous for a deletion of *SPO20* was constructed (see Materials and Methods). This deletion replaces the coding region for amino acids 41–268 of Spo20p with the *URA3* gene, presumably




Figure 2. *SPO20* encodes a second yeast SNAP-25 homologue. Alignment of domains in Sec9p, Spo20p, and mouse SNAP-25. Amino acid numbers are indicated adjacent to hatched areas, and numbers in parentheses are total amino acids. These sequence data are available from GenBank/EMBL/DDBJ under accession numbers Z49211 (*SPO20*), L34336 (*SEC9*), and M22012 (*SNAP-25*).

creating a null allele. The deletion strain, AN67, was examined for secretory phenotypes. No defect was found either in the secretion of invertase or in the sorting of carboxypeptidase Y to the vacuole (Neiman, A.M., unpublished observations); however, AN67 failed to sporulate. Specifically, DAPI staining indicated that the *spo20ΔURA3* mutant strain proceeded through meiosis but failed to form spores, similar to the *sec1*, *sec4*, and *sec8* mutants (Table II).

To examine the *SPO20* gene product further, an epitope-tagged allele was constructed by fusing a sequence encoding three copies of the influenza hemagglutinin epitope (HA) to the 3' end of the coding region. This fusion gene partially complemented the spore formation defect of the *spo20ΔURA3* mutant (see Materials and Methods). To examine the expression of *SPO20*, AN67 carrying the *SPO20-HA* fusion gene under the *SPO20* promoter on a high copy plasmid was sporulated, and cells were removed at various times after transfer to sporulation medium. Extracts were made from these cells, and the proteins were analyzed by Western blot using anti-HA antibodies. Consistent with the lack of any obvious phenotype of the *spo20ΔURA3* mutant in vegetative cells, the protein was not detectable in vegetative cells and was first detected between 3 and 6 h after shift to sporulation medium (Fig. 3). The induction of Spo20p is coincident with or just before the appearance of tetranucleate cells and thus corresponds to the time of the second meiotic division and the onset of prospore membrane formation (Fig. 3). Northern analysis of transcripts from a meiotic time course using *SPO20* as a probe indicates that the transcript is seen only in sporulating cells (Conrad, M., personal communication). Therefore, induction of Spo20p seems to be regulated at the level of transcription.

spo20 Mutants Display a Defect in the Capture of Nuclei by the Prospore Membrane

A simple model based on the sporulation phenotypes of *sec9* and *spo20* is that Sec9p and Spo20p have separate functions. Sec9p is a component of a t-SNARE specifically required for vesicle fusion at the plasma membrane, and Spo20p could play the analogous role in a prospore membrane-specific t-SNARE complex. The remaining fusion machinery (e.g., Sec1p, Sec4p, and Sec8p) may be used for both membranes. If so, then during sporulation *spo20* mutants should have the same cytological phenotype as *sec4* mutants; namely, no prospore membranes should be formed. To examine the *spo20ΔURA3* phenotype more closely, strain AN67 was sporulated, and at various times samples were removed and examined by electron microscop

Surprisingly, the *spo20ΔURA3* mutant forms prospore membranes. Examination of a large number of cells revealed that the *spo20* diploid does display a defect in prospore membrane growth, but a much more subtle one than in *sec4* mutants. At early stages in prospore membrane development in the *spo20* strain, the membranes appear normal and, as in wild type, are juxtaposed to the outer plaque of the spindle pole body (Fig. 4, A–C). In wild-type cells, as the membrane grows it surrounds a lobe of the nucleus (Fig. 4 D). However, in the *spo20* strain, the connection between prospore membrane and nucleus seems to be lost. Prospore membranes are found in the cytoplasm, no longer adjacent to the nucleus (Fig. 4, E and F). These detached prospore membranes continue to grow and eventually fuse with themselves without enclosing a daughter nucleus, thus forming anucleate, immature spores (Fig. 4 H). In some cases, these anucleate spores appear to initiate spore wall development (Fig. 4 I), although no mature spore walls were ever seen.

Spo20p and *Sec9p* Have Partially Overlapping Functions

This cytological analysis demonstrated that, although abnormal, the prospore membrane does form in *spo20* mutants. Analysis of *sec9-4/sec9-4* diploids in the electron microscope showed that spores form normally (data not shown). Taken together, these two results suggest either that fusion of vesicles with the prospore membrane can occur without the function of a SNAP-25 homologue or that




Figure 3. Time course of Spo20p induction. At the times indicated, samples of strain AN67 carrying pRS425-SPO20HA were removed from sporulation medium, and extracts were made and analyzed by Western blot using anti-HA antibodies. Equal cell equivalents were loaded in each lane. Marks at left indicate the position of molecular size standards. The fraction of cells at each time point (at least 200 cells counted) that were tetranucleate, as determined by DAPI staining, is shown below each lane. The predicted molecular mass for the Spo20-HA fusion protein is 49,800 D.




Figure 4. *spo20* mutants fail to package daughter nuclei into spores. (A, D, and G) Isogenic *SPO20* (NH144) or (B, C, E, F, H, and I) *spo20ΔURA3/spo20ΔURA3* (AN67) cells at different stages of prospore membrane formation: early (A–C); middle (D–F); or after closure (G–I). (A and B) The gap in the nuclear envelope underneath the outer plaque is the position of the spindle pole body. N, nucleus; OP, outer plaque; PrM, prospore membrane; SW, spore wall. Bars: (A, B, and I) 200 nm; (C–H) 500 nm.

SEC9 and *SPO20* have overlapping functions. In the latter case, one would expect that prospore membrane formation in the *spo20ΔURA3* diploid would be dependent on *SEC9* function. To test this possibility, a *spo20ΔURA3/spo20ΔURA3 sec9-4/sec9-4* strain (AN80) was constructed. This strain was sporulated at 34°C to inactivate *SEC9* and examined in the electron microscope. The sporulation defect in the double mutant resembled the *sec4-8/sec4-8* mutant in that no prospore membranes were observed (Fig. 5). It seemed possible that the cytological differences seen between AN67 (*spo20ΔURA3/spo20ΔURA3*) and AN80 (*sec9-4/sec9-4 spo20ΔURA3/spo20ΔURA3*) were due to differences in strain background and not the presence of the *sec9-4* mutation. To examine this possibility, a *sec9-4* allele in AN80 was converted to wild type by transformation with a linear DNA fragment carrying the wild-type *SEC9* gene and direct selection for growth at 36°C. The resulting strain (*sec9-4/SEC9 spo20ΔURA3/spo20ΔURA3*) was analyzed by electron microscopy and displayed the same prospore membrane defect described above for AN67 (data

not shown). Thus, the lack of prospore membrane development is a direct consequence of losing both Sec9p and Spo20p function. These results demonstrate that Sec9p plays a role in prospore membrane growth and provide additional evidence that the late acting *SEC* genes are required for fusion of vesicles with the prospore membrane.

To further examine the functional overlap between Sec9p and Spo20p, we tested whether overexpression of one gene could rescue a mutation in the other. Expression of *SPO20* from the *GAL* promoter failed to rescue the *sec9-4* temperature-sensitive phenotype (data not shown). Similarly, expression of *SEC9* from a high copy plasmid could not rescue the *spo20* sporulation defect (data not shown). Thus, though the two proteins have some functional overlap, they cannot completely substitute for each other.

The Cell Surface Protein Gas1p Localizes to the Growing Prospore Membrane

If the prospore membrane is formed by the redirection of




Figure 5. Prospore membranes are absent in a *spo20/spo20 sec9/sec9* double mutant. Cells of strain AN80 (*spo20ΔURA3/spo20ΔURA3sec9-4/sec9-4*) at times corresponding to middle (A) and postclosure stages (B) of prospore membrane formation. N, nucleus; V, vacuole. Cells were sporulated at the restrictive temperature for *sec9-4* (34°C). Bars, 500 nm.

secretory vesicles from the plasma membrane to an intracellular site, then the prospore membrane may also contain the usual cargo of those vesicles. One might expect, for example, to find plasma membrane proteins localized to the growing prospore membrane. To test this hypothesis we have examined the localization of the GPI-anchored cell surface protein Gas1p (Nuoffer et al., 1991) during sporulation (Fig. 6). In tetranucleate cells, Gas1p appears to be localized in an internal membrane that surrounds the nucleus (Fig. 6 A) consistent with a prospore membrane localization. This localization of Gas1p is disrupted in a *sec1-1* mutant strain (data not shown), indicating that this internal membrane is the prospore membrane.

The localization of the secreted chitinase Cts1p (Kuranda and Robbins, 1987) was also examined in sporulating cells. Though the signal is fainter than for Gas1p, Cts1p displays an intracellular staining pattern similar to Gas1p in tetranucleate cells (Fig. 6, C and D, arrowheads). This staining may represent localization to the lumen of the prospore membrane.

Discussion

This report describes evidence for a role of the late-acting *SEC* genes in formation of the prospore membrane. Several late-acting *SEC* genes were examined. *SEC4* and, in a *spo20* background, *SEC9* are required for the formation of this intracellular compartment. *sec1* and *sec8* mutants also fail to sporulate at restrictive temperature, presumably because of a similar defect in prospore membrane formation. These are the first genes shown to be specifically required for formation of the prospore membrane.

In vegetative cells, these *SEC* gene products are required solely for the fusion of post-Golgi vesicles with the plasma membrane. Therefore, the simplest interpretation of these results is that these gene products are required during sporulation for fusion of vesicles to form the prospore membrane. Furthermore, the vesicles that fuse to form the prospore membrane are likely the same ones regulated by late-acting *SEC* genes in vegetative cells, i.e.,




Figure 6. Gas1p and Cts1p localization in sporulating cells. (A) Indirect immunofluorescence of Gas1p in sporulating cells of strain NH144. (B) DAPI staining of cells in A. (C) Indirect immunofluorescence of Cts1p in sporulating cells of strain NH144. (D) DAPI staining of cells in C. Arrowheads (C and D) indicate tetranucleate cells.

secretory vesicles. Consistent with this interpretation, the cell surface protein Gas1p and the secreted protein Cts1p are localized to the prospore membrane compartment in sporulating cells. Thus, trafficking to the prospore membrane defines a new, post-Golgi branch of the yeast secretory pathway (Fig. 7).

During vegetative growth, proteins sorted at the TGN can be packaged into vesicles destined for either the vacuole or the plasma membrane. Only the latter class of vesicles is dependent on Sec4p function for fusion with their target membrane. In sporulating cells, the Sec4p-dependent vesicles could potentially fuse with either the plasma membrane or the prospore membrane (Fig. 7). The presence of two potential target membranes raises the possibility that secretory vesicles are differentially targeted to these two membranes during sporulation.

Recent studies have shown that at least two different classes of secretory vesicle can be distinguished in yeast on the basis of their density (Harsay and Bretscher, 1995). Though all of these vesicles require late-acting *SEC* genes to fuse with the plasma membrane, the different classes contain different cargo molecules. The results presented here do not, therefore, indicate whether all secretion is directed into the prospore membrane or if only a subset of secretory vesicles are directed there. Related to the question of which vesicles are delivered to the prospore membrane is the question of how secretory vesicles are targeted to this intracellular compartment. In vegetative yeast, fusion of secretory vesicles occurs only at particular sites on the cell surface (Winsor and Scheibel, 1997). The restriction of secretion to specified regions of the plasma membrane is mirrored in the polarized organization of the actin cyto-




Figure 7. A sporulation-specific branch of the yeast secretory pathway. Routes for vesicular traffic out of the TGN in vegetative and sporulating cells are shown. Thick circles indicate vesicles dependent on *SEC4* function for fusion with their target membrane. In both situations, *SEC4*-independent vesicles (*thin circles*) are targeted to the vacuole. “?” indicates that it is unclear whether trafficking to the plasma membrane continues in cells developing a prospore membrane (see text).

skeleton, and mutations that disrupt the cytoskeleton are often associated with a delocalization of secretion (Winsor and Scheibel, 1997). Presumably, the targeting of secretory vesicles to the interior of the cell is accompanied by changes in cell polarity and the cytoskeleton. An illustration of this is the redistribution of septins (Fares et al., 1996). In vegetative cells, septins form a ring of 10-nm filaments around the mother/bud neck, whereas in sporulating cells the septins are found in a band (presumably) underlying the growing prospore membrane. It will be of interest to examine a variety of mutations in genes affecting cell polarity and the cytoskeleton to determine their effects on sporulation.

Distinct Functions for *Sec9p* and *Spo20p*

SEC9 and *SPO20* are partially redundant in that the presence of one or the other of these gene products is required for formation of a prospore membrane. However, the single mutations differ in their effects. Mutations in *sec9* have no effect on spore formation, whereas no spores form in *spo20* mutants because the growing prospore membranes lose their attachment to the nucleus and subsequently fail to capture daughter nuclei. One possible explanation for this difference is that *Sec9p* and *Spo20p* are completely redundant but in the absence of *Spo20p*, the level of *Sec9p* during sporulation is insufficient to support proper spore formation. However, overexpression of *SEC9* cannot rescue the *spo20* sporulation defect. Therefore, *Spo20p* appears to have a function, which it does not share with *Sec9p*, that is required to maintain the position of the prospore membrane adjacent to the nucleus.

Similarly, ectopic expression of *SPO20* in vegetative cells cannot suppress the temperature-sensitive phenotype of a *sec9-4* mutant, suggesting that *Sec9p* has some essential function that *Spo20p* cannot provide. This result is surprising given that *Spo20p* can replace *Sec9p* during sporulation. It may be that there are additional sporulation-specific proteins required for *Spo20p* to function in vesicle fusion. Alternatively, *Sec9p* and *Spo20p* may mediate the fusion of overlapping but distinct sets of vesicles so that

neither protein can completely compensate for a defect in the other.

Secretion-driven Cell Division in Other Systems

Spore formation is a process in which cell division occurs not by a classical cytokinetic mechanism but rather by the growth of new plasma membranes, discontinuous from the mother cell plasma membrane, to give rise to daughter cells within the cytoplasm of the mother cell. This form of cell division has features in common with “nonclassical” division events in a number of systems. In *Drosophila*, for example, during both cellularization of the syncytial blastoderm and spermatogenesis, polynucleate cells are divided into mononucleate cells by processes that involve the coalescence of vesicles to form new plasma membranes (Tokuyasu et al., 1972; Loncar and Singer, 1995). Similarly, in higher plants cell plate formation occurs first by the coalescence of vesicles in the interior of the cell between the daughter nuclei (Wick, 1991; Staehelin and Hepler, 1996). The vesicles then fuse to form a flattened sheet, which then grows outward by vesicle addition until it fuses with the mother cell plasma membrane to complete division.

Perhaps the most striking parallel to ascospore morphogenesis is also found in plants: the formation of the generative cell during pollen development (McCormick, 1993). The four haploid products of the male meiosis are termed microspores. Each of these microspores then undergoes a mitotic division. One of the daughter nuclei from this division will remain in the mother cell cytoplasm and is termed the vegetative nucleus. The second is enclosed in a double membrane within the mother cell cytoplasm. This smaller cell is termed the generative cell. The generative cell will eventually undergo a mitotic division to give rise to two sperm cells. Together, these three cells (two sperm and one vegetative) form the mature pollen grain. The double membrane around the generative cell is formed by the coalescence of vesicles around the generative cell nucleus (Gimenez-Martin et al., 1969; Ledbetter and Porter, 1970) and appears directly analogous to the formation of spores in *S. cerevisiae*. A cell wall even forms between the two membranes surrounding the generative cell (Angold, 1968; Heslop-Harrison, 1968), similarly to spore wall maturation. It will be of interest to learn if these different events in plants, yeast, and metazoans share some common molecular mechanisms, such as the use of alternative SNAP-25 proteins.

I thank N. Hollingsworth, R. Sternglanz, P. Pryciak (University of Massachusetts, Worcester, MA), N. Dean, and an anonymous reviewer for comments on the manuscript; N. Hollingsworth, C. Kaiser (Massachusetts Institute of Technology, Cambridge, MA), and D. Terbush (Yale University, New Haven, CT) for strains; and M. Conrad (Oklahoma Medical Research Foundation) for communication of results before publication. I am grateful to N. Dean for antibodies and to G. Rudomen for able assistance with the electron microscopy. I am also indebted to N. Hollingsworth, N. Dean, B. Aronson, and R. Sternglanz for discussion, advice, and encouragement throughout the course of this project.

A.M. Neiman was supported by the Cancer Research Fund of the Damon Runyon-Walter Winchell Foundation, DRG-1276 and National Institutes of Health grant GM-28220 to R. Sternglanz.

Received for publication 21 May 1997 and in revised form 3 November 1997.

References

- Aalto, M.K., H. Ronne, and S. Keranen. 1993. Yeast syntaxins Sso1p and Sso2p belong to a family of related membrane proteins that function in vesicular transport. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:4095-4104.
- Angold, R.E. 1968. The formation of the generative cell in the pollen grain of *Emdmyion non-scriptus* (L.). *J. Cell Sci.* 3:573-578.
- Brennwald, P., B. Kearns, K. Champion, S. Keranen, V. Bankaitis, and P. Novick. 1994. Sec9 is a SNAP-25-like component of a yeast SNARE complex that may be the effector of Sec4 function in exocytosis. *Cell.* 79:245-258.
- Briza, P., M. Breitenbach, A. Ellinger, and J. Segall. 1990. Isolation of two developmentally regulated genes involved in spore wall maturation in *Saccharomyces cerevisiae*. *Genes Dev.* 4:1775-1789.
- Byers, B. 1981. Cytology of the yeast life cycle. In *The Molecular Biology of the Yeast Saccharomyces: Life Cycle and Inheritance*. J.N. Strathern, E.W. Jones, and J.R. Broach, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY. 59-96.
- Esposito, R.E., and S. Klapholz. 1981. Meiosis and ascospore development. In *The Molecular Biology of the Yeast Saccharomyces: Life Cycle and Inheritance*. J.N. Strathern, E.W. Jones, and J.R. Broach, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY. 211-287.
- Fares, H., L. Goetsch, and J.R. Pringle. 1996. Identification of a developmentally regulated septin and involvement of the septins in spore formation in *Saccharomyces cerevisiae*. *J. Cell Biol.* 132:399-411.
- Friesen, H., R. Lunz, S. Doyle, and J. Segall. 1994. Mutation of the *SPS1*-encoded protein kinase of *Saccharomyces cerevisiae* leads to defects in transcription and morphology during spore formation. *Genes Dev.* 8:2162-2175.
- Gimenez-Martin, G., M.C. Risueno, and J.F. Lopez-Saez. 1969. Generative cell envelope in pollen grains as a secretion system, a postulate. *Protoplasma.* 67: 223-235.
- Harsay, E., and A. Bretscher. 1995. Parallel secretory pathways to the cell surface in yeast. *J. Cell Biol.* 131:297-310.
- Heslop-Harrison, J. 1968. Synchronous pollen mitosis and the formation of generative cell in massulata orchids. *J. Cell Sci.* 3:457-466.
- Krisak, L., R. Strich, R.S. Winters, J.P. Hall, M.J. Mallory, D. Kreitzer, R.S. Tuan, and E. Winter. 1994. SMK1, a developmentally regulated MAP kinase, is required for spore wall assembly in *Saccharomyces cerevisiae*. *Genes Dev.* 8:2151-2161.
- Kupiec, M., B. Byers, R.E. Esposito, and A.P. Mitchell. 1997. Meiosis and sporulation in *Saccharomyces cerevisiae*. In *The Molecular and Cellular Biology of the Yeast Saccharomyces Cell Cycle and Cell Biology*. J.R. Pringle, J.R. Broach, and E.W. Jones, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY. 889-1036.
- Kuranda, M.J., and P.W. Robbins. 1987. Cloning and heterologous expression of glycosidase genes from *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA.* 84:2585-2589.
- Ledbetter, M.C., and K.R. Porter. 1970. Introduction to the Fine Structure of Plant Cells. Springer-Verlag, New York. 270 pp.
- Loncar, D., and S.J. Singer. 1995. Cell membrane formation during the cellularization of the syncytial blastoderm of *Drosophila*. *Proc. Natl. Acad. Sci. USA.* 92:2199-2203.
- Lynn, R.R., and P.T. Magee. 1970. Development of the spore wall during ascospore formation in *Saccharomyces cerevisiae*. *J. Cell Biol.* 44:688-692.
- McCormick, S. 1993. Male gametophyte development. *Plant Cell.* 5:1265-1275.
- Moens, P.B. 1971. Fine structure of ascospore development in the yeast *Saccharomyces cerevisiae*. *Can. J. Microbiol.* 17:507-510.
- Moens, P.B., and E. Rapport. 1971. Spindles, spindle plaques, and meiosis in the yeast *Saccharomyces cerevisiae* (Hansen). *J. Cell Biol.* 50:344-361.
- Moens, P.B., R.E. Esposito, and M.S. Esposito. 1974. Aberrant nuclear behavior at meiosis and anucleate spore formation by sporulation-deficient (SPO) mutants of *Saccharomyces cerevisiae*. *Exp. Cell Res.* 83:166-174.
- Neiman, A.M., V. Mhaiskar, V. Manus, F. Galibert, and N. Dean. 1997. *Saccharomyces cerevisiae HOC1*, a suppressor of *pkc1*, encodes a putative glycosyltransferase. *Genetics.* 145:637-645.
- Novick, P., S. Ferro, and R. Schekman. 1981. Order of events in the yeast secretory pathway. *Cell.* 25:461-469.
- Nuoffer, C., P. Jenö, A. Conzelmann, and H. Riezman. 1991. Determinants for glycosylphospholipid anchoring of the *Saccharomyces cerevisiae* GAS1 protein to the plasma membrane. *Mol. Cell. Biol.* 11:27-37.
- Pelham, H.R.B., K.G. Hardwick, and M.J. Lewis. 1988. Sorting of soluble ER proteins in yeast. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:1757-1762.
- Protopopov, V., B. Govindan, P. Novick, and J.E. Gerst. 1993. Homologues of the synaptobrevin/VAMP family of synaptic vesicle proteins function in the late secretory pathway in *S. cerevisiae*. *Cell.* 74:855-861.
- Rose, K., S.A. Rudge, M.A. Frohman, A.J. Morris, and J. Engebrecht. 1995. Phospholipase D signaling is essential for meiosis. *Proc. Natl. Acad. Sci. USA.* 92:12151-12155.
- Rose, M.D., and G.R. Fink. 1990. *Methods in Yeast Genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 119-187.
- Rothman, J.E. 1994. Mechanisms of intracellular protein transport. *Nature.* 372: 55-63.
- Rothman, J.H., C.P. Hunter, L.A. Valls, and T.H. Stevens. 1986. Overproduction-induced mislocalization of a yeast vacuolar protein allows isolation of its structural gene. *Proc. Natl. Acad. Sci. USA.* 83:3248-3252.
- Rothstein, R.J. 1983. One-step gene disruption in yeast. *Methods Enzymol.* 101: 202-211.
- Salminen, A., and P.J. Novick. 1987. A ras-like protein is required for a post-Golgi event in yeast secretion. *Cell.* 49:527-538.
- Sikorski, R.S., and P. Hieter. 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics.* 122:19-27.
- Söllner, T., S.W. Whiteheart, M. Brunner, H. Erdjument-Bromage, S. Geromanos, P. Tempst, and J.E. Rothman. 1993. SNAP receptors implicated in vesicle targeting and fusion. *Nature.* 362:318-324.
- Søgaard, M., K. Tani, R. Ruby ye, S. Geromanos, P. Tempst, T. Kirchhausen, J.E. Rothman, and T. Söllner. 1994. A Rab protein is required for the assembly of SNARE complexes in the docking of transport vesicles. *Cell.* 78:937-948.
- Staehelein, L.A., and K.P. Hepler. 1996. Cytokinesis in higher plants. *Cell.* 84: 821-824.
- TerBush, D.R., and P. Novick. 1995. Sec6, Sec8, and Sec15 are components of a multisubunit complex which localizes to small bud tips in *Saccharomyces cerevisiae*. *J. Cell Biol.* 130:299-312.
- Tokuyasu, K.T., W.J. Peacock, and R.W. Hardy. 1972. Dynamics of spermiogenesis in *Drosophila melanogaster*. I. Individualization process. *Z. Zellforsch.* 124:479-506.
- Tu, J., W. Song, and M. Carlson. 1996. Protein phosphatase type 1 interacts with proteins required for meiosis and other cellular processes in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 16:4199-4206.
- Wick, S.M. 1991. Spatial aspects of cytokinesis in plant cells. *Curr. Opin. Cell Biol.* 3:253-260.
- Winsor, B., and E. Schiebel. 1997. An overview of the *Saccharomyces cerevisiae* microtubule and microfilament cytoskeleton. *Yeast.* 13:399-434.

