

# A molecular switch that controls cell spreading and retraction

Panagiotis Flevaris, Aleksandra Stojanovic, Haixia Gong, Athar Chishti, Emily Welch, and Xiaoping Du

Department of Pharmacology, University of Illinois at Chicago, Chicago, IL 60612

Integrin-dependent cell spreading and retraction are required for cell adhesion, migration, and proliferation, and thus are important in thrombosis, wound repair, immunity, and cancer development. It remains unknown how integrin outside-in signaling induces and controls these two opposite processes. This study reveals that calpain cleavage of integrin  $\beta_3$  at Tyr<sup>759</sup> switches the functional outcome of integrin signaling from cell spreading to retraction. Expression of a calpain cleavage-resistant  $\beta_3$  mutant in Chinese hamster ovary cells causes defective clot retraction and RhoA-mediated retraction signaling but

enhances cell spreading. Conversely, a calpain-cleaved form of  $\beta_3$  fails to mediate cell spreading, but inhibition of the RhoA signaling pathway corrects this defect. Importantly, the calpain-cleaved  $\beta_3$  fails to bind c-Src, which is required for integrin-induced cell spreading, and this requirement of  $\beta_3$ -associated c-Src results from its inhibition of RhoA-dependent contractile signals. Thus, calpain cleavage of  $\beta_3$  at Tyr<sup>759</sup> relieves c-Src-mediated RhoA inhibition, activating the RhoA pathway that confines cell spreading and causes cell retraction.

## Introduction

Integrins are a family of heterodimeric cell adhesion receptors, which not only mediate cell adhesion to extracellular matrix proteins but also transmit signals that are vital for anchorage-dependent cell survival, proliferation, and motility (Hynes, 2002). Thus, integrins play pivotal roles in physiological processes such as wound healing, immune responses, and hemostasis. Aberrant integrin signaling is also centrally involved in the development of human diseases such as thrombosis, cancer, and autoimmune diseases and is the target of many therapeutic agents (Hynes, 2002; Shattil and Newman, 2004). Therefore, understanding the molecular events in the transduction of integrin signals is important in our understanding of many biological processes and has the potential to reveal novel intervention for diseases.

Integrin signaling is bidirectional in that ligand binding function of integrins requires conformational change in the extracellular ligand binding domain induced by intracellular signals (inside-out signaling), and ligand binding transduces outside-in signals that mediate cellular responses (Hynes, 2002; Vinogradova et al., 2002; Ginsberg et al., 2005; Wegener et al., 2007). An early functional outcome of integrin outside-in signaling is cell spreading, which is characterized by the formation of filopodia and lamellipodia and represents the outward movement of

cell membrane, polymerizing actin, and cytoskeletal complexes at the leading edge (Hall, 2005). Integrin signaling later induces cell retraction, which is the inward movement of the cell membrane and cytoskeletal complexes, often at the rear end of cells. Coordinated spreading and retraction allows cell migration and in blood platelets facilitates stable platelet adhesion, thrombus formation, and consolidation. It has been shown that small GTP binding proteins Rac and cdc42 are involved in signaling mechanisms of cell spreading and that RhoA-mediated signaling is important in cell retraction (Hall, 2005). However, it remains unclear how integrin signaling initiates and temporally regulates these two seemingly opposing cellular responses; this is a fundamental question in cell biology. In this study, we have discovered that calpain cleavage of  $\beta_3$  at Tyr<sup>759</sup> serves as a molecular switch that changes the outcome of the integrin outside-in signals from mediating cell spreading to promoting cell retraction. Furthermore, the switch from cell spreading to retraction is mediated by calpain cleavage of the c-Src binding site at the integrin C terminus, which relieves the c-Src-dependent inhibition of RhoA, and thus facilitates integrin-mediated, RhoA-dependent contractile signaling.

## Results

### A calpain cleavage-resistant $\beta_3$ mutant

We have previously reported that the calcium-dependent protease calpain cleaves the cytoplasmic domain of the integrin  $\beta_3$

Correspondence to Xiaoping Du: xdu@uic.edu

Abbreviations used in this paper: RBD, Rho binding domain; ROCK, Rho-dependent kinase; SFK, Src family kinase; WT, wild type.

subunit mainly at Y<sup>759</sup>, thus removing the C-terminal RGT<sup>762</sup> sequence (Du et al., 1995; Xi et al., 2003). Calpain cleavage of  $\beta_3$  is inhibited by tyrosine phosphorylation at Y<sup>759</sup> of the  $\beta_3$  cytoplasmic domain (Xi et al., 2006). To understand the physiological role of calpain cleavage of  $\beta_3$ , we developed a calpain-resistant mutant of  $\beta_3$  by replacing R<sup>760</sup> with a negatively charged glutamic acid (R760E; Fig. 1 A). This mutation is intended to mimic the effect of phosphorylation at the adjacent Y<sup>759</sup> by inhibiting calpain cleavage without perturbing the functionally critical NITY motif. The R760E mutant was stably expressed in a CHO cell line (Gu et al., 1999) together with the wild-type (WT) integrin  $\alpha_{IIb}\beta_3$  subunit. To determine whether the R760E mutant confers resistance to calpain-mediated proteolysis, lysates from cells expressing WT integrin  $\alpha_{IIb}\beta_3$  and the R760E mutant were incubated with or without purified  $\mu$ -calpain and immunoblotted with Ab762, an antibody recognizing the intact  $\beta_3$  C-terminal TYRGT<sup>762</sup> sequence, or Ab759, an antibody that recognizes the  $\beta_3$  TNITY<sup>759</sup> sequence only when  $\beta_3$  is

cleaved by calpain at Y<sup>759</sup> (Du et al., 1995; Xi et al., 2003). The specificities of these antibodies have been previously characterized in Du et al. (1995) and Xi et al. (2003). In particular, we have shown that the calpain cleavage-specific antibody Ab759 only reacts with the  $\beta_3$  that is truncated at the C-terminal side of Y<sup>759</sup> and fails to recognize  $\beta_3$  that is not cleaved (WT) or truncated at different sites, and that Ab762 reacts only with WT  $\beta_3$  that is not cleaved and fails to react with  $\beta_3$  cleaved at Y<sup>759</sup> or other sites (Xi et al., 2003). MAb 15, recognizing the  $\beta_3$  extracellular domain (which is not disturbed by calpain; Du et al., 1995), serves as loading control (Fig. 1 B). WT  $\beta_3$  shows a large increase in reactivity with Ab759 after treatment with  $\mu$ -calpain. This increase is mirrored by the loss of reactivity with Ab762, indicating loss of intact C terminus. In contrast, calpain-treated R760E mutant  $\beta_3$  showed no decrease in Ab762 reactivity and dramatically diminished reactivity with Ab759. The relatively low reactivity of Ab762 with the R760E mutant compared with WT  $\beta_3$  is caused by R<sup>760</sup> to E mutation but is not




Figure 1. **A calpain cleavage-resistant mutation of  $\beta_3$ : R760E.** (A) WT  $\beta_3$  cytoplasmic domain sequence and the R760E mutation. (B) Cells expressing WT and R760E mutant  $\alpha_{IIb}\beta_3$  were solubilized, treated with or without purified  $\mu$ -calpain, and immunoblotted with anti- $\beta_3$  C-terminal antibody Ab762, calpain cleavage-indicator antibody Ab759, and the anti- $\beta_3$  extracellular domain antibody mAb 15. Note that although R760E showed lower reactivity with Ab762 because of mutation in the epitope (TYR<sup>760</sup>GT), calpain caused minimal loss of Ab762 reactivity or gain of Ab759 reactivity, which indicates calpain resistance. (C) Analysis of stable  $\alpha_{IIb}\beta_3$  expression in cells expressing WT and R760E mutant integrins, using the monoclonal anti- $\alpha_{IIb}\beta_3$  antibody D57. (D) WT and R760E cells spreading on fibrinogen were permeabilized and stained with mAb 15 (green) and purified Ab759 (red). Ab759 staining of R760E cells was reduced compared with WT  $\beta_3$ . Bar, 10  $\mu$ m.

caused by reduced R760E expression, as indicated by the reaction with mAb 15. These results demonstrate that R760E mutation is calpain cleavage resistant (Fig. 1 B). Levels of stable surface expression of the R760E mutant and WT  $\alpha_{IIb}\beta_3$  were also analyzed by flow cytometry to ensure comparable expression after repeated cell sorting with an anti- $\alpha_{IIb}\beta_3$  monoclonal antibody, D57 (Fig. 1 C). We showed previously that the  $\beta_3$  cytoplasmic domain is cleaved by endogenous calpain in platelets and

integrin-expressing CHO cells adherent to fibrinogen (Xi et al., 2003). To test whether R760E is resistant to endogenous calpain during cell adhesion, CHO cells containing either WT cells or R760E mutant  $\beta_3$  were allowed to adhere and spread on immobilized fibrinogen and were subsequently immunolabeled with affinity-purified cleavage-specific antibody Ab759 to detect integrin cleavage and with mAb 15 to mark  $\beta_3$ . WT  $\beta_3$  was cleaved by calpain (Fig. 1 D). In contrast, the R760E mutant




Figure 2. **Effects of cleavage-resistant  $\beta_3$  mutation, R760E, and calpain inhibition on clot retraction and  $\beta_3$  cleavage.** (A) Fibrinogen-depleted plasma mixed with WT or R760E cells was induced to coagulate with 2 U/ml thrombin at 37°C for the indicated times, and then photographed. (B) Quantification of clot size at various time points in A. Error bars are the SD from three separate experiments. (C) Clots containing WT or R760E cells were fixed at 1 h, cryosectioned, stained with purified Ab759 (red) or mAb 15 (green), and observed by confocal microscopy. Note the reduced Ab759 staining of R760E cells. Bars, 10  $\mu$ m. (D) Clot retraction by WT cells preincubated with either 5  $\mu$ M of calpain inhibitor MDL 28170 or vehicle control. (E) Quantification of clot sizes in D (mean  $\pm$  SD). (F) Washed platelets from WT and  $\mu$ -calpain knockout mice were incubated in an aggregometer with or without 0.1 U/ml thrombin for 7 min.  $\beta_3$  cleavage at Y<sup>759</sup>, which was detected using cleavage-specific antibody Ab759, was enhanced in the WT compared with the calpain knockout. Ab8053 recognizes  $\beta_3$  extracellular domain and serves as a loading control.

showed marked reduction in calpain cleavage (Fig. 1 D). Thus, the R760E mutation renders  $\beta_3$  resistant to cleavage by calpain in cells adherent to fibrinogen.

#### **The effects of calpain resistance on $\beta_3$ -mediated clot retraction and cell spreading**

Fibrin clot retraction, mediated by platelets and  $\beta_3$  integrin-expressing cells, requires  $\beta_3$ -dependent outside-in signaling and cell retraction (Chen et al., 1994; Law et al., 1996). To determine whether and how the calpain-resistant mutation of  $\beta_3$  affects integrin signaling, we compared the ability of the WT  $\beta_3$ - and R760E-expressing cells to mediate clot retraction (Fig. 2, A and B). Clot retraction mediated by R760E cells was significantly decreased ( $P < 0.001$ ). Consistent with this result, calpain cleaved WT  $\beta_3$  at Y<sup>759</sup> during clot retraction, and this cleavage was impeded in the R760E mutant cells (Fig. 2 C). It was previously reported that calpain I knockout platelets showed reduced ability to mediate clot retraction (Azam et al., 2001). Similarly, a calpain inhibitor, MDL28170, also reduced clot retraction mediated by  $\beta_3$  integrin-expressing CHO cells (Fig. 2, D and E). However, the calpain substrate responsible for this effect of calpain was unclear. Thus, our data represent a new finding that calpain cleavage of  $\beta_3$  stimulates clot retraction. This finding is consistent with the data obtained in platelets. Although massive integrin cleavage induced by calcium ionophore or high concentrations of thrombin was not significantly affected in calpain I<sup>-/-</sup> platelets in a previous study (possibly because of the involvement of Calpain II; Azam et al., 2001; Pfaff et al., 1999), we show in Fig. 2 F that calpain cleavage of  $\beta_3$  at Y<sup>759</sup> was considerably reduced in calpain I<sup>-/-</sup> platelets when the cleavage was induced by a lower concentration of thrombin (0.1 U/ml), which is consistent with the finding that the calpain I<sup>-/-</sup> mice were defective in clot retraction when clot retraction was stimulated with a low concentration of thrombin (Azam et al., 2001).

To determine whether the calpain-resistant  $\beta_3$  mutant affects integrin-dependent cell spreading, cells expressing WT or R760E mutant  $\alpha_{IIb}\beta_3$  were allowed to adhere to fibrinogen, and spreading kinetics of these cells were observed (Fig. 3, A and B). R760E mutation significantly enhanced and accelerated cell spreading compared with WT  $\alpha_{IIb}\beta_3$  ( $P < 0.0001$ ), indicating that calpain cleavage of  $\beta_3$  negatively regulates cell spreading. This result is consistent with the data that cells expressing a deletion mutant of  $\beta_3$  mimicking calpain cleavage at Y<sup>759</sup> ( $\Delta 759$  cells) are defective in integrin-dependent cell spreading (Fig. 3 E). Thus, calpain cleavage of  $\beta_3$  negatively regulates the integrin outside-in signals that promote cell spreading while it stimulates the integrin outside-in signaling that activates cellular retractile machinery.

#### **The inhibitory effect of calpain cleavage of $\beta_3$ on cell spreading is mediated by activation of RhoA-dependent cell retraction**


The increase in clot retraction and inhibition of cell spreading by calpain cleavage of  $\beta_3$  can result either from cleavage-induced activation of cell retractile function or cleavage-induced inhibition of

cell spreading signaling. To differentiate these two possibilities, we sought to inhibit the cell retractile signaling mechanisms. Previous studies demonstrate that stress fiber formation in cells adherent to integrin ligands requires activation of RhoA (Jaffe and Hall, 2003) and Rho-dependent kinases (ROCK), which stimulate myosin light chain phosphorylation (Amano et al., 1996; Kimura et al., 1996) and thus retractile function of the actomyosin complex (Giuliano et al., 1992). Indeed, the ROCK inhibitors Y-27632 or H-1152, when preincubated with WT  $\beta_3$ -expressing CHO cells (Fig. 3 C) and platelets (Fig. 3 D), impeded clot retraction, indicating that the activation of the RhoA-ROCK signaling pathway is important in integrin-dependent cell retraction. To determine whether this pathway is responsible for calpain cleavage-dependent reversal of the direction of cell membrane movement, we studied the effects of ROCK inhibitors on cell spreading of WT  $\beta_3$ -expressing or calpain cleavage-mimicking  $\beta_3$ -expressing cells. The CHO cells expressing the calpain-cleaved form of  $\beta_3$  ( $\Delta 759$ ) were defective in spreading on fibrinogen, and this defect was completely corrected by the ROCK inhibitors (Fig. 3 E). Spreading of CHO cells expressing WT  $\alpha_{IIb}\beta_3$  was not significantly enhanced ( $P > 0.05$ ) by low concentrations of ROCK inhibitors (Fig. 3, E and F). However, higher concentrations of either H-1152 or Y-27632 accelerate cell spreading in WT cells (Fig. 3, G and H). Interestingly, enhancement of spreading by cells expressing cleavage-resistant integrin mutant R760E was not observed even with higher-dose ROCK inhibition (Fig. 3, G and H), suggesting that the inhibitory effect of Rho-kinase signaling on cell spreading is already diminished when  $\beta_3$  is resistant to calpain cleavage. These results suggest that cleavage of  $\beta_3$  at Y<sup>759</sup> caused activation of the RhoA-ROCK signaling pathway and cell retraction, thus preventing cell spreading.

#### **Calpain cleavage of $\beta_3$ at Y<sup>759</sup> induces RhoA activation**

RhoA is activated by GTP binding and inactivated when bound GTP is hydrolyzed to GDP (Jaffe and Hall, 2003). The Rho binding domain (RBD) of the RhoA effector protein rhotekin specifically binds GTP-bound RhoA and can therefore be used to indicate RhoA activation (Ren and Schwartz, 2000). To determine whether calpain cleavage of  $\beta_3$  stimulates RhoA activation, cells were allowed to spread on fibrinogen-coated surfaces, and then were fixed, permeabilized, and incubated with fluorescent GST-RBD fusion protein (GST-RBD-555). GST-RBD-555 strongly reacted with the WT integrin-expressing cells spreading on fibrinogen (Fig. 4). GST-RBD-555 binding was specific because it was competitively inhibited by unlabeled GST-RBD (Fig. 4 A) and by preincubation of cells with cell-permeable C3 transferase (Fig. 5) but was unaffected in the presence of 20 mM glutathione (Fig. 4 A). The specificity of GST-RBD-555 is further supported by its colocalization with the stain of a monoclonal antibody that recognizes RhoA (Fig. 4 B). Thus, by using this novel in situ RhoA activation assay, we show that RhoA is activated after WT integrin-mediated cell adhesion and spreading on fibrinogen. In contrast to WT  $\beta_3$ , cells expressing the calpain-resistant  $\beta_3$  mutant R760E display reduced reactivity with GST-RBD-555, indicating that the resistance to calpain cleavage by R760E mutation and increased






**Figure 3. Effects of cleavage-resistant and -mimicking  $\beta_3$  mutations and ROCK inhibitors on cell spreading and clot retraction.** (A) Indicated cell lines were incubated on fibrinogen-coated chamber slides at 37°C and photographed at various times. Images at 20 min are shown. (B) Sizes of spread cells were quantified as percentage of original sizes using Image J software. Histogram shows the mean  $\pm$  SEM. (C and D) 5  $\mu$ M each of ROCK inhibitors Y-27632 and H-1152 inhibited clot retraction in platelets (D) and in CHO cells expressing WT  $\beta_3$  (C). (E) Spreading of WT cells and cells expressing a calpain cleavage-mimicking  $\beta_3$  mutant ( $\Delta$ 759) on fibrinogen for 1 h. Defective spreading of  $\Delta$ 759 cells was corrected by 2  $\mu$ M each of ROCK inhibitors H-1152 and Y-27632. (F) Quantification of cell spreading from cells in E, using image J software. Histogram shows the mean  $\pm$  SEM. (G) 5  $\mu$ M each of ROCK inhibitors H-1152 and Y-27632 enhance spreading in WT cells but not in cleavage-resistant R760E mutants at 20 min. (H) Quantification of cell spreading from cells in G, using image J software. Histogram shows the mean  $\pm$  SEM.

cell spreading in R760E cells are associated with inhibition of integrin-induced RhoA activation. Conversely,  $\Delta$ 759 cells, which are defective in spreading on fibrinogen, showed robust reac-

tivity with GST-RBD-555, indicating that the calpain-cleaved form of  $\beta_3$  induces RhoA activation without requiring cell spreading. Collectively, these data demonstrate that cleavage of  $\beta_3$  at Y<sup>759</sup>

Figure 4. **Calpain cleavage of  $\beta_3$  stimulates RhoA activation.** (A) WT, R760E, and  $\Delta 759$  cells spreading on fibrinogen for 1 h were fixed and allowed to react with Alexa Fluor 555-labeled rhotekin (GST-RBD-555 [red]) to indicate RhoA activation. To estimate nonspecific binding, fluorescent rhotekin binding was also performed in the presence of unlabeled rhotekin (NC) or 20 mM glutathione (glut). MAb 15 (green) serves as a marker for cell margins and integrin  $\beta_3$  expression. (B) Activated RhoA, indicated by GST-RBD-555 (red), colocalizes with anti-RhoA (green). Bars, 10  $\mu\text{m}$ .



switches on the RhoA-ROCK signaling pathway, leading to cell retraction and inhibition of cell spreading.

**Effects of calpain cleavage-resistant or -mimicking mutations on c-Src binding to  $\beta_3$**

It has been reported that a member of the Src family of protein kinases (SFK), c-Src, interacts with the  $\beta_3$  cytoplasmic domain

and that this interaction requires the YRGT<sup>762</sup> sequence in the C terminus of  $\beta_3$  (Arias-Salgado et al., 2005a,b). To determine whether c-Src interaction with  $\beta_3$  is affected by the R760E mutation or calpain cleavage-mimicking truncation of  $\beta_3$ , WT and mutant  $\beta_3$ , stably expressed in CHO cells, were immunoprecipitated with Ab8053, an anti- $\beta_3$  antibody, and precipitates were immunoblotted for  $\beta_3$ -associated c-Src (Fig. 6). We found that




Figure 5. **Inhibition of GST-RBD binding to active RhoA by cell-permeable C3 transferase.** Cells expressing WT  $\alpha_{IIb}\beta_3$  were preincubated with or without 5  $\mu\text{g}/\text{ml}$  of cell-permeable C3 transferase and were allowed to spread on fibrinogen-coated surfaces for 1 h. Cells were stained with GST-RBD-555 (red) to detect active RhoA and with mAb 15 (green) to mark integrin surface expression. Bars, 10  $\mu\text{M}$ .

c-Src coimmunoprecipitated with both WT and R760E mutant forms of  $\beta_3$  but failed to interact with the cleavage-mimicking mutant  $\Delta 759$  (Fig. 6). These data are consistent with previous studies (Arias-Salgado et al., 2005a,b), indicating that cleavage of the  $\beta_3$  C-terminal RGT sequence disrupts c-Src binding to  $\beta_3$ .

**Calpain cleavage of  $\beta_3$  at Y<sup>759</sup> relieves the inhibitory effect of c-Src on RhoA activation and thus promotes cell retraction**

It has been shown that c-Src plays an important role in cell spreading (Kaplan et al., 1995) and that c-Src interaction with  $\beta_3$  is important for this function (Arias-Salgado et al., 2003; Shattil, 2005). Therefore, we examined whether the loss of  $\beta_3$ -associated c-Src function is responsible for the activation of cell retraction and inhibition of cell spreading induced by calpain cleavage at Y<sup>759</sup>. PP2, an SFK inhibitor, abolished WT  $\beta_3$ -mediated cell spreading on fibrinogen (Fig. 7, A and D). The inhibitory effect of PP2 bears striking resemblance to the deficiency in cell spreading in  $\Delta 759$  cells. More importantly, the inhibitory effect of PP2 was reversed by ROCK inhibitors H-1152 and Y-27632, indicating that the RhoA-ROCK-mediated retractile signaling is required for the inhibitory effect of this SFK inhibitor on cell spreading (Fig. 7, A and D), which is similar to the requirement for ROCK in  $\beta_3$  cleavage-mediated inhibition of cell spreading. To exclude the possible nonspecific effect of pharmacological inhibitors, WT  $\beta_3$ -expressing cells were transfected with cDNA, encoding a dominant-negative mutant of c-Src (K295R/Y527F; Mukhopadhyay et al., 1995), and cotransfected with 1/6 quantity of GFP cDNA to indicate successful transfection. The control cells expressing transfected GFP exhibited normal cell spreading on fibrinogen (Fig. 7, B and E). In contrast, the cells transfected with the dominant-negative c-Src showed a defect in cell spreading similar to that caused by PP2 (Fig. 7, B and D). This inhibitory effect of the dominant-negative c-Src was reversed when cells were cotransfected with an equal concentration of dominant-negative RhoA (N19-RhoA; Fig. 7, B and D; Qiu et al., 1995). The data obtained with cDNA transfection is thus consistent with the data using inhibitors of SFK and ROCK, indicating that integrin-dependent cell spreading occurs independent of c-Src when RhoA-mediated retractile signaling is disabled.

Because integrin-mediated c-Src activation inhibits RhoA but does not affect cdc42 and Rac functions (Arthur et al., 2000), our data indicate that c-Src promotes integrin-dependent cell spreading by its inhibition of RhoA-mediated retractile signals. Furthermore, the SFK inhibitor PP2 corrected the defective RhoA activation in calpain cleavage-resistant R760E cells, restoring RhoA activation and inhibiting the spreading of R760E cells (Fig. 7, C and F), which is similar to the phenotype observed in  $\Delta 759$  cells expressing the calpain-cleaved form of  $\beta_3$ . These results indicate that without calpain cleavage of  $\beta_3$ ,  $\beta_3$ -dependent c-Src activity inhibits RhoA activation and thus promotes cell spreading. Conversely, inhibition of SFK with PP2 dramatically enhanced clot retraction (Fig. 7 G), further demonstrating that SFK inhibits cell retractile machinery. Therefore, cleavage of  $\beta_3$  at Y<sup>759</sup> relieves the inhibitory effect of c-Src on RhoA activation, switches on RhoA-ROCK-dependent cell retraction, and thus changes the functional outcome of integrin signaling from cell spreading to cell retraction.






Figure 6. **Coimmunoprecipitation of c-Src with WT  $\beta_3$  and  $\beta_3$  mutants.** (A)  $\beta_3$  immunoprecipitates from CHO cells expressing WT, R760E mutant, and  $\Delta 759$  mutant integrins were probed on Western blots with antibodies against  $\beta_3$  (mAb 15) or c-Src. c-Src binding to the integrin cytoplasmic domain is abolished in the calpain cleavage-mimicking mutant  $\Delta 759$  but not in the calpain cleavage-resistant mutant R760E. (B) Lysates were also immunoblotted with antibody to c-Src.





**Figure 7. The effect of calpain cleavage of  $\beta_3$  on c-Src-dependent inhibition of RhoA-mediated cell retraction.** (A) WT cells incubated with or without 10  $\mu$ M PP2 or PP2 plus 4  $\mu$ M of ROCK inhibitors H-1152 or Y-27632 were allowed to spread on fibrinogen for 1 h. Inhibition of cell spreading by PP2 is reversed by ROCK inhibitors. Bar, 10  $\mu$ m. (B) Spreading of WT cells transfected with GFP alone, a dominant-negative c-Src (DN c-Src) plus GFP, N19-RhoA plus GFP, or N19-RhoA and dominant-negative c-Src plus GFP. Note that N19-RhoA reversed the inhibitory effect of dominant-negative c-Src. (C) R760E cells are allowed to adhere to fibrinogen in the presence of PP2 or vehicle control DMSO and are stained with GST-RBD-555 (red) and mAb 15 (green). Defective RhoA activation in R760E cells is corrected by PP2. Bars, 10  $\mu$ m. (D) Quantitation of cell sizes in A using Image J (mean  $\pm$  SEM from four random fields). (E) Quantitation of cell sizes in B using Image J (mean  $\pm$  SEM from three experiments). (F) Stain of GST-RBD-555 in cells from C was quantified using LSM 5 software (mean  $\pm$  SEM from three experiments). (G) 10  $\mu$ M of SFK inhibitor PP2 accelerates clot retraction in platelets.





**Figure 8. The effect of c-Src activation on the integrin-cleavage-dependent inhibition of cell spreading.** (A) WT, R760E, and  $\Delta 759$  cells were either in suspension or allowed to spread on fibrinogen for 30 min, and then solubilized and analyzed by Western blot with antibodies specific for c-Src or Src phosphorylated at Y<sup>416</sup> as an indicator of Src activation. Note that c-Src activation is similar in all three cell lines. (B)  $\Delta 759$  cells were transfected with either GFP or GFP plus a constitutively active mutant of c-Src, E378G. Nonadherent transfected cells were solubilized and immunoblotted with antibodies against total c-Src, Y<sup>416</sup>-phosphorylated c-Src, and  $\beta_3$  (Ab8053, to verify equal loading). (C)  $\Delta 759$  cells were transfected with GFP alone or GFP plus a constitutively active mutant of c-Src, E378G. The transfected cells were allowed to spread on fibrinogen-coated surfaces. Note that active c-Src cannot reverse the spreading defect in  $\Delta 759$  cells. (D) Quantitation of cell sizes in C using Image J software (mean  $\pm$  SEM from three experiments).

### Cleavage of $\beta_3$ at Y<sup>759</sup> diminishes c-Src-dependent cell spreading by dissociating c-Src from $\beta_3$ but not by inhibiting overall c-Src activation

Previous studies suggested that activity of c-Src associated with  $\beta_3$  is increased compared with that of c-Src that is not associated with  $\beta_3$  (Arias-Salgado et al., 2005a). To investigate the possibility that cleavage of  $\beta_3$  at Y<sup>759</sup> regulates integrin-dependent overall c-Src activation in adherent cells, we measured the amount of active c-Src in CHO cells expressing WT and calpain cleavage-resistant (R760E) and -mimicking ( $\Delta 759$ ) mutants of  $\beta_3$ . All three cell lines were similarly capable of inducing considerable c-Src phosphorylation at Y<sup>416</sup> (an indicator of c-Src activation) when allowed to adhere and spread onto fibrinogen-coated surfaces (Fig. 8 A), suggesting that deletion of the c-Src binding site in  $\beta_3$  did not affect the overall integrin-mediated c-Src activation. Furthermore, if c-Src activity alone was sufficient to mediate the inhibition of RhoA, constitutively active c-Src would be able to induce integrin-mediated cell spreading. Thus,  $\Delta 759$  cells were transfected with cDNA encoding a constitutively active mutant of c-Src (E378G) and cotransfected with 1/6 quantity of GFP cDNA to indicate successful transfection. Increased levels of total c-Src and expression of constitutively active c-Src (Y<sup>416</sup> phosphorylated) in E378G c-Src-transfected  $\Delta 759$  cells in suspension were verified by Western blot (Fig. 8 B).

The E378G c-Src-transfected  $\Delta 759$  cells are not different from the GFP-expressing control  $\Delta 759$  cells in that both spread poorly on fibrinogen (Fig. 8, C and D). These data suggest that the inhibitory effect of  $\beta_3$  cleavage on cell spreading is not because of its effect on overall c-Src activation but is associated with dissociation of c-Src (and therefore c-Src activity) from the  $\beta_3$  cytoplasmic domain. Thus, only the  $\beta_3$ -associated c-Src activity is responsible for mediating RhoA inhibition and promoting cell spreading. Collectively, our results strongly suggest that cleavage of  $\beta_3$  at Y<sup>759</sup> dissociates c-Src from the  $\beta_3$  C-terminal domain and thus relieves the inhibitory effect of  $\beta_3$ -associated c-Src on RhoA activation, stimulating the RhoA-ROCK retractile signals, which switches the functional outcome of integrin signaling from mediating cell spreading to promoting cell retraction.

## Discussion

In this study, we have discovered that  $\beta_3$  cleavage by calpain serves as a molecular switch that changes the outcome of integrin outside-in signaling from mediating cell spreading to promoting cell retraction. The switch in the outcome of integrin signaling from cell spreading to retraction is mediated by  $\beta_3$  cleavage-dependent activation of the RhoA retractile signaling pathway, and  $\beta_3$  cleavage-dependent activation of the RhoA signaling pathway is mediated by cleavage-dependent relief of

the inhibitory effects of  $\beta_3$ -associated c-Src on the RhoA signaling pathway. Thus, our study also reveals that the requirement of SFK in integrin-dependent cell spreading is caused by its inhibition of the RhoA-dependent retractile function of the cell. These new findings represent major conceptual advances in our understanding of how integrin signaling controls and regulates the direction of cell membrane movement.


The data supporting our conclusions in this study are mainly obtained using the reconstituted integrin signaling model in CHO cells expressing WT and calpain cleavage-resistant or -mimicking mutants of  $\beta_3$ . Integrins play critical roles in all mammalian cells that adhere to extracellular matrix. Although different cell types express different members of the integrin family, integrin signaling, particularly outside-in signaling, is generally conserved in adherent cells, including CHO cells. In the past 20 yr, many important findings in  $\beta_3$  integrin function and signaling have been discovered using the CHO cell model expressing the WT and mutant  $\beta_3$  integrins, including the finding of the important role of talin in integrin activation (O'Toole et al., 1990; Calderwood et al., 1999; Tadokoro et al., 2003; Xi et al., 2003). Furthermore, the results obtained in CHO cells have been consistently verified in native  $\beta_3$ -expressing cells, such as platelets (Li et al., 2001, 2003a, 2006; Tadokoro et al., 2003; Zou et al., 2007), indicating the validity of the CHO cell model. Importantly, the data obtained in CHO cell models in this study are consistent with the role of calpain I in clot retraction and in  $\beta_3$  cleavage as indicated in experiments using human and mouse platelets. Nevertheless, it is recognized that CHO cells are not totally identical to native  $\beta_3$ -expressing cells such as platelets, and thus the results obtained with the CHO cell model will need to be further verified in platelets and other  $\beta_3$ -expressing cells in the future.

We have previously shown that the cytoplasmic domain of  $\beta_3$  is a calpain substrate and that the cleavage occurs mainly at Y<sup>759</sup>, removing the C-terminal RGT sequence (Du et al., 1995; Xi et al., 2003). However, although calpain is activated after calcium elevation is induced by ligand binding to the integrin, very little cleavage of  $\beta_3$  occurs early during cell spreading because of the protection of integrin from calpain cleavage by tyrosine phosphorylation of the  $\beta_3$  cytoplasmic domain, especially at the periphery of a cell (Xi et al., 2006). In contrast, at a later time during cell spreading when Y<sup>759</sup> of  $\beta_3$  becomes dephosphorylated, substantial  $\beta_3$  cleavage occurs (Xi et al., 2006). Also, calpain cleavage of  $\beta_3$  is readily detectable both during clot retraction in platelets and in the  $\beta_3$ -expressing CHO cells. The kinetics of integrin cleavage are consistent with our data, indicating a role for calpain in the transition between cell spreading and cell retraction in both CHO cells and in platelets. In fact, the data obtained with cells expressing the calpain-resistant and calpain cleavage-mimicking mutants clearly reveal that cleavage of  $\beta_3$  at Y<sup>759</sup> generates the change in integrin signaling from mediating cell spreading to promoting cell retraction. Thus, we have discovered a calpain-dependent molecular switch that is regulated temporally and topographically by tyrosine phosphorylation of  $\beta_3$  and that controls the direction of membrane and cytoskeleton movement induced by integrin outside-in signaling. It is important to note that the cleavage-mimicking mutant of

$\beta_3$  creates an experimental condition where 100% of  $\beta_3$  cytoplasmic domain is cleaved at Y<sup>759</sup>, which serves to clearly reveal the cleavage-induced changes in functional outcome. In platelets, >22% of  $\beta_3$  is cleaved at Y<sup>759</sup> within an hour after platelet activation (Xi et al., 2003), which coincides with the occurrence of marked clot retraction. However, in more delicate situations, such as during cell spreading and migration, local and dynamic calpain activation signals are likely to cause localized cleavage of a small percentage of  $\beta_3$  molecules and thus dynamically control the local movement of membranes and cytoskeleton.

The  $\beta_3$  cleavage-mediated inhibition of cell spreading and promotion of cell retraction is not caused by its inhibition of cell spreading mechanisms but is mediated by the integrin cleavage-dependent activation of the RhoA retractile signaling pathway. Previous studies suggest that Rac and cdc42 play important roles in mediating cell spreading (filopodia and lamellipodia formation) and that RhoA is increasingly activated in spread cells (Hall, 2005). RhoA activates ROCK, which phosphorylates and inactivates myosin light chain phosphatase, and enhances myosin light chain phosphorylation and retractile function of the actomyosin complex (Giuliano et al., 1992; Amano et al., 1996; Kimura et al., 1996), inducing cell retraction. Cleavage of  $\beta_3$  at Y<sup>759</sup> inhibits integrin-mediated cell spreading (Fig. 3; Xi et al., 2003). If the inhibitory effects of  $\beta_3$  cleavage on cell spreading should result from its inhibition of Rac- and cdc42-mediated cell spreading signaling, this inhibitory effect should not be corrected by inhibition of RhoA pathway. Therefore, our data that  $\beta_3$  cleavage-dependent inhibition of cell spreading is corrected by ROCK inhibitors indicate that calpain cleavage of  $\beta_3$  activates RhoA-dependent cell retraction, which subsequently inhibits the opposite movement of cell membrane resulting in inhibition of cell spreading.

Previous studies indicate that members of SFK are required for integrin-mediated signaling that leads to cell spreading (Shattil, 2005). The mechanism responsible for this role of SFK is not clear but is believed to be an early step in cell spreading signaling. However, we show that the inhibitory effect of an SFK inhibitor and a dominant-negative c-Src mutant on  $\beta_3$ -mediated cell spreading is reversed by inhibition of the RhoA signaling pathway, indicating that cell spreading is independent of c-Src when RhoA signaling is abrogated. These data therefore provide the first direct evidence that the role of SFK in promoting integrin-dependent cell spreading is not that it is required for cdc42- and Rac-dependent cell spreading signaling, but rather is its inhibition of RhoA-dependent retractile signaling and thus cell retraction, which confines the outward movement of the cell membrane and cytoskeleton. Our data are consistent with the previous studies, which show that SFK inhibits RhoA activity but has no effect on the activity of cdc42 and Rac (Arthur et al., 2000). Our data are also consistent with the finding that RhoA inactivation by p190RhoGAP promotes membrane protrusion and polarity (Arthur and Burridge, 2001). More importantly, we show that cleavage of  $\beta_3$  at Y<sup>759</sup> abolishes c-Src binding to  $\beta_3$  but does not significantly affect overall c-Src activation in cells spread on fibrinogen, and that the RhoA-dependent inhibition of cell spreading by  $\beta_3$  cleavage at Y<sup>759</sup> is not reversed by expression of a constitutively active Src. These data suggest



**Figure 9. A new calpain-dependent switch that controls cell spreading and retraction.** This schematic illustrates the calpain-dependent switch that controls the direction of integrin signaling. In this model, ligand binding to integrins induces not only Rac- and cdc42-dependent cell spreading signals, but also phosphorylation of the cytoplasmic domain of  $\beta_3$  and elevation of intracellular calcium levels that activate calpain. Tyrosine phosphorylation at Y<sup>759</sup> of  $\beta_3$  protects the c-Src binding site in the C-terminal domain of  $\beta_3$  from calpain cleavage. The  $\beta_3$ -associated c-Src facilitates cell spreading by inhibiting RhoA-dependent cell retraction signals. In spread cells, Y<sup>759</sup> of  $\beta_3$  becomes dephosphorylated, allowing  $\beta_3$  cleavage by calpain. Removal of the c-Src binding site in  $\beta_3$  by calpain relieves the local inhibitory effect of c-Src on the RhoA-ROCK signaling pathway, activates RhoA-dependent cell retraction, and thus switches the outcome of local integrin signaling from mediating cell spreading to retraction. However, the possible involvement of additional pathways is not totally excluded.

that whereas overall intracellular c-Src activation during cell adhesion does not require c-Src association with  $\beta_3$ , active c-Src molecules dissociated from  $\beta_3$  are not capable of inhibiting the RhoA signaling pathway and promoting cell spreading on  $\beta_3$  ligands, possibly because of the lack of  $\beta_3$  scaffold. However, the calpain cleavage-resistant mutation in  $\beta_3$  inhibits RhoA activation in a Src-dependent manner. Thus, our results suggest that only the  $\beta_3$ -associated c-Src is responsible for the inhibition of the RhoA pathway and facilitation of cell spreading. Conversely, our results suggest that calpain cleavage of  $\beta_3$ , which abolishes Src binding to  $\beta_3$  (Fig. 6; Arias-Salgado et al., 2005b), relieves the inhibitory effects of  $\beta_3$ -associated c-Src on the RhoA pathway, activates RhoA, and thus switches the outcome of integrin signaling from cell spreading to retraction.

Interestingly,  $\beta_3$  cleavage-dependent relief of c-Src-mediated RhoA inhibition and activation of RhoA-dependent cell retraction is associated with the cleavage of only a fraction of  $\beta_3$  molecules (Fig. 2; Xi et al., 2003) and cannot be prevented by the presence of uncleaved  $\beta_3$  molecules in the same cell. This apparently dominant effect of  $\beta_3$  cleavage suggests that c-Src activity at one location in a cell does not cross-inhibit RhoA activation induced by calpain cleavage of  $\beta_3$  at a different location and is consistent with the knowledge that spreading (such as filopodia and lamellipodia) and retraction can occur simultaneously at different locations within a cell (such as during cell migration). The requirement of c-Src association with  $\beta_3$  for the c-Src-mediated RhoA inhibition provides a mechanism for the localized role of c-Src in promoting cell spreading. Further studies are required to understand how  $\beta_3$ -associated c-Src activity

locally regulates RhoA and whether additional mechanisms are needed for RhoA activation when  $\beta_3$  cleavage by calpain relieves the inhibitory effect of  $\beta_3$ -associated c-Src.

Based on data from this and other studies, we propose a new model of integrin signaling that incorporates a phosphorylation- and protease-mediated mechanism regulating integrin-mediated cell membrane movements. In this model (Fig. 9), initial integrin signaling activates protein tyrosine kinases, which induce phosphorylation of the cytoplasmic domain of  $\beta_3$  (Law et al., 1996, 1999), and elevates intracellular calcium levels, which activate calpain (Fox et al., 1993). Tyrosine phosphorylation at Y<sup>759</sup> of  $\beta_3$  in the early phase of cell spreading protects the c-Src binding site in the C-terminal domain of  $\beta_3$  from calpain cleavage (Xi et al., 2006). The interaction of c-Src with  $\beta_3$  promotes activation of  $\beta_3$ -associated c-Src by a Csk- (Oberfell et al., 2002) and protein tyrosine phosphatase IB-dependent mechanism (Arias-Salgado et al., 2005a) and/or allows c-Src-dependent signaling in the vicinity of the  $\beta_3$  C terminus, which facilitates cell spreading by inhibiting RhoA-dependent cell retraction. In spread cells, Y<sup>759</sup> of  $\beta_3$  becomes dephosphorylated (Xi et al., 2006), allowing  $\beta_3$  cleavage by activated calpain, which removes the C-terminal RGT sequence of  $\beta_3$  important in c-Src binding. Integrin cleavage by calpain thus relieves the local inhibitory effect of  $\beta_3$ -associated c-Src on the RhoA-ROCK signaling pathway and locally activates RhoA-dependent cell retractile signals (Xi et al., 2003; Arias-Salgado et al., 2005b). Thus, calpain cleavage of  $\beta_3$  switches the direction of integrin outside-in signaling from mediating spreading to promoting cell retraction.



## Materials and methods

### Reagents

Plasma containing 0.38% sodium citrate and 40  $\mu\text{g}/\text{ml}$  aprotinin (Sigma-Aldrich) was depleted of fibronectin as previously reported in Chen et al. (1994). Ab762 and Ab759 were previously described in Du et al. (1995) and Xi et al. (2003). mAb 15, D57, and Ab8053 were provided by M. Ginsberg (University of California, San Diego, La Jolla, CA; O'Toole et al., 1990). Phosphospecific anti-Src Y<sup>416</sup> antibody was obtained from Cell Signaling; anti-RhoA and cell permeable C3 transferase were obtained from Cytoskeleton, Inc.; Sudan black was obtained from Sigma-Aldrich; Alexa Fluor 594-conjugated goat anti-rabbit IgG and Alexa Fluor 488-conjugated goat anti-mouse IgG were obtained from Invitrogen; anti- $\nu$ -Src monoclonal antibody, SFK inhibitor PP2, ROCK inhibitors H-1152 and Y-27632, and calpain inhibitor MDL 28170 were obtained from Calbiochem; and an anti-c-Src polyclonal antibody was obtained from Santa Cruz Biotechnology, Inc. GST-RBD cDNA were a gift from T. Kozasa (University of Illinois, Chicago, IL). GST-RBD fusion protein was purified as previously described in Ren and Schwartz (2000). Purified protein was fluorescently labeled with an Alexa Fluor 555 microscale protein labeling kit (Invitrogen).

### Construction of R760E mutant

The  $\beta_3$  R760E mutation was performed using PCR. The forward primer has the sequence of AGAGCTTAAGGACAC. The reverse primers (CTCATTAAGTCCCTCGTAGGTGATATTGG) contain an XhoI digestion site, stop codon, and the 18-nucleotide C-terminal  $\beta_3$  sequence coding the intended R760E mutation. PCR products were digested with AspI and XhoI and ligated into a  $\beta_3$  cDNA construct in a modified cDM8 vector containing only one XhoI site at the 3' end of the multiple cloning site. The construct was verified by DNA sequencing.

### Cell lines and transfection

CHO cells stably expressing WT human  $\alpha_{IIb}\beta_3$  (2b3a), human glycoprotein Ib-IX (GPIb-IX; 1b9), or both  $\alpha_{IIb}\beta_3$  and GPIb-IX (123) were previously described in Gu et al. (1999) and Xi et al. (2003). CHO cells expressing GPIb-IX and a truncation mutant  $\beta_3$ , mimicking calpain cleavage at Y<sup>759</sup>, and complexed with WT  $\alpha_{IIb}$  ( $\Delta 759$ ) were also described previously in Gu et al. (1999) and Xi et al. (2003). DNA transfection was performed using Lipofectamine 2000 (Invitrogen). R760E  $\beta_3$  mutant cDNA was co-transfected with WT  $\alpha_{IIb}$  and pcDNA 3.1/Hyg plasmid at a ratio of 5:5:1 into 1b9 cells. Complex formation between  $\alpha_{IIb}$  and  $\beta_3$  subunits was verified by flow cytometry using antibody D57 against  $\alpha_{IIb}\beta_3$  complex. Stable R760E mutant cell lines were obtained by antibiotic selection and repeated cell sorting using D57 as previously described (Gu et al., 1999). To ensure comparable expression, R760E and  $\Delta 759$  cells were sorted using the expression level of the stable WT  $\alpha_{IIb}\beta_3$ -expressing (123) cells as a gate (Gu et al., 1999; Xi et al., 2003). For transient transfections, cells stably coexpressing WT  $\alpha_{IIb}\beta_3$  were transfected with either 2  $\mu\text{g}$  GFP, 2  $\mu\text{g}$  GFP plus 12  $\mu\text{g}$  of dominant-negative c-Src (Mukhopadhyay et al., 1995), 2  $\mu\text{g}$  GFP plus 12  $\mu\text{g}$  of dominant-negative N19-RhoA (Qiu et al., 1995), or 2  $\mu\text{g}$  GFP plus 12  $\mu\text{g}$  of both dominant-negative c-Src and N19-RhoA, using Lipofectamine 2000. In some experiments, cells were transiently transfected with either 2  $\mu\text{g}$  GFP or 2  $\mu\text{g}$  GFP plus 12  $\mu\text{g}$  of constitutively active c-Src (E378G). Cells were analyzed 24 h after transfection.

### Calpain cleavage of $\beta_3$ in CHO cells and mouse platelets

$10^7$  CHO cells were directly solubilized in buffer containing 2% Triton X-100, 0.15 M NaCl, 1 mM CaCl<sub>2</sub>, and 0.02 M Tris, pH 7.4, in the presence or absence of 10 mM EDTA. Cell lysates were incubated for 40 min at 37°C with or without 1  $\mu\text{g}$  of purified  $\mu$ -calpain. SDS-PAGE sample buffer containing 10 mM EDTA, 1 mM PMSF, and 0.2 mM E64 was added to stop reactions. For experiments using mice, 6–8-wk-old mice of either sex were anesthetized by intraperitoneal injection of pentobarbital and blood was drawn from the inferior vena cava. Blood from 5–6 (calpain I<sup>+/+</sup> and calpain I<sup>-/-</sup>; Azam et al., 2001) mice was pooled or platelets were isolated by differential centrifugation as described in Li et al. (2003b).  $3 \times 10^8/\text{ml}$  washed platelets were resuspended in modified Tyrode's buffer (Du et al., 1991) and incubated at 22°C for 2 h before use. Platelets were stimulated with 0.1 U/ml thrombin in a platelet aggregometer at 37°C for 7 min, and then solubilized in SDS-PAGE sample buffer. All samples were analyzed by SDS-PAGE using 4–15% gradient gels and were immunoblotted using various antibodies. Enhanced chemiluminescence (GE Healthcare) was used for visualization of antibody reactions.

### Immunofluorescence and confocal microscopy

Chamber slides (Lab-Tek; Nunc) were precoated with 10  $\mu\text{g}/\text{ml}$  fibrinogen and blocked with 5% BSA in PBS. 50  $\mu\text{l}$  CHO cells (expressing WT or mutant integrins;  $5 \times 10^5/\text{ml}$ ) in Tyrode's buffer was added to the wells and incubated at 37°C for 1 h. After washing, cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100, 0.1 M Tris, 10 mM EGTA, 0.15 M NaCl, 5 mM MgCl<sub>2</sub>, 1 mM PMSF, 0.1 mM E64, and 1% BSA (permeabilization buffer, pH 7.5). Samples were blocked with 5% BSA, incubated with 10  $\mu\text{g}/\text{ml}$  mAb 15 and 0.5  $\mu\text{g}/\text{ml}$  of purified Ab759, and, after washing, stained with fluorescently labeled secondary antibodies. To detect RhoA activation, cells were fixed in 4% paraformaldehyde in PBS with 10 mM MgCl<sub>2</sub> and incubated with 0.7  $\mu\text{g}/\text{ml}$  mAb 15 and Alexa Fluor 555-labeled GST-RBD in permeabilization buffer containing 10 mM MgCl<sub>2</sub>. After further staining with Alexa Fluor 488-conjugated goat anti-mouse IgG and washes, the antibody and GST-RBD staining were detected using a confocal microscope (Carl Zeiss MicroImaging, Inc.). In some experiments, cells were also preincubated with or without 5  $\mu\text{g}/\text{ml}$  of cell-permeable C3 transferase to inhibit RhoA, according to the manufacturer's instructions, and then stained with GST-RBD-555. Surface area in spreading cells was estimated by measuring total pixel number per cell in digitally recorded images using Image J software (National Institutes of Health). Quantitation of 555-nm fluorescence in R760E cells was determined by using the area tool in LSM 5 software (Carl Zeiss MicroImaging, Inc.) and expressed as total pixel number. Statistical significance was determined using a *t* test.

### Clot retraction

400  $\mu\text{l}$  of indicated CHO cell lines ( $4 \times 10^6/\text{ml}$ ), in complete DME in the presence or absence of 5  $\mu\text{M}$  ROCK inhibitors H-1152 or Y-27632 or calpain inhibitor MDL 28170, was mixed with 100  $\mu\text{l}$  of plasma, 2 U/ml thrombin, and 5 mM CaCl<sub>2</sub> in a partially Sigmacote-treated glass cuvette (Chrono-Log). For platelet samples, citrated platelet-rich plasma was mixed with 0.2 U/ml thrombin. The clots were allowed to retract at 37°C and were photographed at various times. The two-dimensional sizes of retracted clots on photographs were quantified using Image J software and were expressed as clot size. Statistical significance was determined using a *t* test.

### Confocal microscopy analysis of clot sections

Clots were fixed with 4% paraformaldehyde, embedded in 50% Tissue-Tek OCT compound (Sakura Finetek), placed in  $25 \times 20 \times 5$ -mm Tissue-Tek Cryomolds (Sakura Finetek), and snap frozen in liquid nitrogen. Frozen sections were made using a cryostat microtome (Microm HM 505 E; Carl Zeiss MicroImaging, Inc.). Slides were then washed and quenched with 5% Sudan black (wt/vol) in 70% ethanol. Sections were permeabilized with 0.1% Triton X-100, 0.1 M Tris, 10 mM EGTA, 0.15 M NaCl, 5 mM MgCl<sub>2</sub>, 1 mM PMSF, 0.1 mM E64, and 1% BSA, pH 7.5 (permeabilization buffer), blocked with 5% BSA, and immunolabeled with 10  $\mu\text{g}/\text{ml}$  mAb 15 and 0.5  $\mu\text{g}/\text{ml}$  of purified Ab 759. After washing, cells were stained with Alexa Fluor 594-conjugated anti-rabbit IgG and 488-conjugated anti-mouse IgG. Data was collected using a confocal microscope.

### Spreading assays

CHO cells expressing WT, R760E mutant, and  $\Delta 759$  mutant integrins ( $100 \mu\text{l}/\text{well}$ ,  $1.5 \times 10^5/\text{ml}$ ) in complete DME were incubated in fibrinogen-coated chamber slides at 37°C for 20, 40, and 60 min. At each time, cells were immediately rinsed and fixed with 4% paraformaldehyde. For assays using PP2 and ROCK inhibitors, cells were preincubated at 22°C for 30 min with or without 10  $\mu\text{M}$  PP2 and/or 4  $\mu\text{M}$  H-1152 or Y-27632. Cells were allowed to spread for 1 h at 37°C, rinsed, and fixed with 4% paraformaldehyde. Images were captured using a microscope (DM IRB; Leica). The size of the cells from four random fields for each time and cell line was determined using Image J software. Relative increase in cell size was calculated by dividing the mean size of spread cells by the mean size of suspended cells from each cell line. Statistical significance was determined using a *t* test.

### Immunoprecipitation

Cells were lysed in lysis buffer containing 1% NP-40, 150 mM NaCl, 50 mM Tris, pH 7.4, 1 mM sodium orthovanadate, 1 mM NaF, and Complete protease inhibitor mixture (Roche Molecular Biochemicals). Lysates were incubated with either rabbit anti- $\beta_3$  or an equal amount of rabbit IgG and subsequently with protein A-Sepharose (GE Healthcare). Beads were washed four times with lysis buffer, analyzed by SDS-PAGE using 4–15% gradient gels, and immunoblotted with mAb 15 for  $\beta_3$  and monoclonal anti-Src (Calbiochem).



### Immunoblot analysis of c-Src activation in cells adherent on fibrinogen-coated surfaces

10-cm polystyrene dishes (Fisher Scientific) were coated with 10  $\mu\text{g}/\text{ml}$  fibrinogen and blocked with 5% BSA in PBS. Equal suspensions containing  $5 \times 10^6$  CHO cells stably expressing WT, R760E mutant, or  $\Delta 759$  mutant integrins in modified Tyrode's buffer were either kept in suspension or added to the dishes and incubated at 37°C for 30 min. Plates were rinsed three times with PBS and cells were subsequently lysed with the buffer containing 1% NP-40, 150 mM NaCl, 50 mM Tris, pH 7.4, 1 mM sodium orthovanadate, 1 mM NaF, and Complete protease inhibitor mixture. Lysates were analyzed by SDS-PAGE using 4–15% gradient gels, and then immunoblotted with anti-phospho-Src Y<sup>416</sup> and anti-c-Src.

We thank Dr. Sanford Shattil for critical reading of the manuscript, Dr. Tohru Kozasa and Nicole Hajjick for helping with the Rho activation experiments, and Dr. Michael Burleigh for helping with histochemistry.

This work is supported by grants from the National Institutes of Health/National Heart, Lung, and Blood Institute (HL080264, HL062350, and HL068819; to X. Du).

Submitted: 28 March 2007

Accepted: 1 October 2007

## References

- Amano, M., M. Ito, K. Kimura, Y. Fukata, K. Chihara, T. Nakano, Y. Matsuura, and K. Kaibuchi. 1996. Phosphorylation and activation of myosin by Rho-associated kinase (Rho-kinase). *J. Biol. Chem.* 271:20246–20249.
- Arias-Salgado, E.G., S. Lizano, S. Sarkar, J.S. Brugge, M.H. Ginsberg, and S.J. Shattil. 2003. Src kinase activation by direct interaction with the integrin beta cytoplasmic domain. *Proc. Natl. Acad. Sci. USA.* 100:13298–13302.
- Arias-Salgado, E.G., F. Haj, C. Dubois, B. Moran, A. Kasirer-Friede, B.C. Furie, B. Furie, B.G. Neel, and S.J. Shattil. 2005a. PTP-1B is an essential positive regulator of platelet integrin signaling. *J. Cell Biol.* 170:837–845.
- Arias-Salgado, E.G., S. Lizano, S.J. Shattil, and M.H. Ginsberg. 2005b. Specification of the direction of adhesive signaling by the integrin beta cytoplasmic domain. *J. Biol. Chem.* 280:29699–29707.
- Arthur, W.T., and K. Burridge. 2001. RhoA inactivation by p190RhoGAP regulates cell spreading and migration by promoting membrane protrusion and polarity. *Mol. Biol. Cell.* 12:2711–2720.
- Arthur, W.T., L.A. Petch, and K. Burridge. 2000. Integrin engagement suppresses RhoA activity via a c-Src-dependent mechanism. *Curr. Biol.* 10:719–722.
- Azam, M., S.S. Andrabi, K.E. Sahr, L. Kamath, A. Kuliopulos, and A.H. Chishti. 2001. Disruption of the mouse mu-calpain gene reveals an essential role in platelet function. *Mol. Cell Biol.* 21:2213–2220.
- Calderwood, D.A., R. Zent, R. Grant, D.J. Rees, R.O. Hynes, and M.H. Ginsberg. 1999. The Talin head domain binds to integrin beta subunit cytoplasmic tails and regulates integrin activation. *J. Biol. Chem.* 274:28071–28074.
- Chen, Y.P., T.E. O'Toole, J. Ylanne, J.P. Rosa, and M.H. Ginsberg. 1994. A point mutation in the integrin beta 3 cytoplasmic domain (S752→P) impairs bidirectional signaling through alpha IIb beta 3 (platelet glycoprotein IIb-IIIa). *Blood.* 84:1857–1865.
- Du, X.P., E.F. Plow, A.L.I. Frelinger, T.E. O'Toole, J.C. Loftus, and M.H. Ginsberg. 1991. Ligands "activate" integrin  $\alpha\text{IIb}\beta 3$  (platelet GPIIb-IIIa). *Cell.* 65:409–416.
- Du, X., T.C. Saïdo, S. Tsubuki, F.E. Indig, M.J. Williams, and M.H. Ginsberg. 1995. Calpain cleavage of the cytoplasmic domain of the integrin  $\beta 3$  subunit. *J. Biol. Chem.* 270:26146–26151.
- Fox, J.E., R.G. Taylor, M. Taffarel, J.K. Boyles, and D.E. Goll. 1993. Evidence that activation of platelet calpain is induced as a consequence of binding of adhesive ligand to the integrin, glycoprotein IIb-IIIa. *J. Cell Biol.* 120:1501–1507.
- Ginsberg, M.H., A. Partridge, and S.J. Shattil. 2005. Integrin regulation. *Curr. Opin. Cell Biol.* 17:509–516.
- Giuliano, K.A., J. Kolega, R.L. DeBiasio, and D.L. Taylor. 1992. Myosin II phosphorylation and the dynamics of stress fibers in serum-deprived and stimulated fibroblasts. *Mol. Biol. Cell.* 3:1037–1048.
- Gu, M., X. Xi, G.D. Englund, M.C. Berndt, and X. Du. 1999. Analysis of the roles of 14-3-3 in the platelet glycoprotein Ib-IX-mediated activation of integrin  $\alpha\text{IIb}\beta 3$  using a reconstituted mammalian cell expression model. *J. Cell Biol.* 147:1085–1096.
- Hall, A. 2005. Rho GTPases and the control of cell behaviour. *Biochem. Soc. Trans.* 33:891–895.
- Hynes, R.O. 2002. Integrins: bidirectional, allosteric signaling machines. *Cell.* 110:673–687.
- Jaffe, A.B., and A. Hall. 2003. Cell biology. Smurfing at the leading edge. *Science.* 302:1690–1691.
- Kaplan, K.B., J.R. Swedlow, D.O. Morgan, and H.E. Varmus. 1995. c-Src enhances the spreading of src<sup>-/-</sup> fibroblasts on fibronectin by a kinase-independent mechanism. *Genes Dev.* 9:1505–1517.
- Kimura, K., M. Ito, M. Amano, K. Chihara, Y. Fukata, M. Nakafuku, B. Yamamori, J. Feng, T. Nakano, K. Okawa, et al. 1996. Regulation of myosin phosphatase by Rho and Rho-associated kinase (Rho-kinase). *Science.* 273:245–248.
- Law, D.A., A.L. Nannizzi, and D.R. Phillips. 1996. Outside-in integrin signal transduction.  $\alpha\text{IIb}\beta 3$ -(GPIIb-IIIa) tyrosine phosphorylation induced by platelet aggregation. *J. Biol. Chem.* 271:10811–10815.
- Law, D.A., F.R. DeGuzman, P. Heiser, K. Ministri-Madrid, N. Killen, and D.R. Phillips. 1999. Integrin cytoplasmic tyrosine motif is required for outside-in  $\alpha\text{IIb}\beta 3$  signalling and platelet function. *Nature.* 401:808–811.
- Li, Z., X. Xi, and X. Du. 2001. A mitogen-activated protein kinase-dependent signaling pathway in the activation of platelet integrin  $\alpha\text{IIb}\beta 3$ . *J. Biol. Chem.* 276:42226–42232.
- Li, Z., X. Xi, M. Gu, R. Feil, R.D. Ye, M. Eigenthaler, F. Hofmann, and X. Du. 2003a. A stimulatory role for cGMP-dependent protein kinase in platelet activation. *Cell.* 112:77–86.
- Li, Z., G. Zhang, G.C. Le Breton, X. Gao, A.B. Malik, and X. Du. 2003b. Two waves of platelet secretion induced by thromboxane A2 receptor and a critical role for phosphoinositide 3-kinases. *J. Biol. Chem.* 278:30725–30731.
- Li, Z., G. Zhang, R. Feil, J. Han, and X. Du. 2006. Sequential activation of p38 and ERK pathways by cGMP-dependent protein kinase leading to activation of the platelet integrin  $\alpha\text{IIb}\beta 3$ . *Blood.* 107:965–972.
- Mukhopadhyay, D., L. Tsiokas, X.M. Zhou, D. Foster, J.S. Brugge, and V.P. Sukhatme. 1995. Hypoxic induction of human vascular endothelial growth factor expression through c-Src activation. *Nature.* 375:577–581.
- O'Toole, T.E., J.C. Loftus, X.P. Du, A.A. Glass, Z.M. Ruggeri, S.J. Shattil, E.F. Plow, and M.H. Ginsberg. 1990. Affinity modulation of the alpha IIb beta 3 integrin (platelet GPIIb-IIIa) is an intrinsic property of the receptor. *Cell Regul.* 1:883–893.
- Obergfell, A., K. Eto, A. Mocsai, C. Buensuceso, S.L. Moores, J.S. Brugge, C.A. Lowell, and S.J. Shattil. 2002. Coordinate interactions of Csk, Src, and Syk kinases with  $\alpha\text{IIb}\beta 3$  initiate integrin signaling to the cytoskeleton. *J. Cell Biol.* 157:265–275.
- Pfaff, M., X. Du, and M.H. Ginsberg. 1999. Calpain cleavage of integrin beta cytoplasmic domains. *FEBS Lett.* 460:17–22.
- Qiu, R.G., J. Chen, F. McCormick, and M. Symons. 1995. A role for Rho in Ras transformation. *Proc. Natl. Acad. Sci. USA.* 92:11781–11785.
- Ren, X.D., and M.A. Schwartz. 2000. Determination of GTP loading on Rho. *Methods Enzymol.* 325:264–272.
- Shattil, S.J. 2005. Integrins and Src: dynamic duo of adhesion signaling. *Trends Cell Biol.* 15:399–403.
- Shattil, S.J., and P.J. Newman. 2004. Integrins: dynamic scaffolds for adhesion and signaling in platelets. *Blood.* 104:1606–1615.
- Tadokoro, S., S.J. Shattil, K. Eto, V. Tai, R.C. Liddington, J.M. de Pereda, M.H. Ginsberg, and D.A. Calderwood. 2003. Talin binding to integrin beta tails: a final common step in integrin activation. *Science.* 302:103–106.
- Vinogradova, O., A. Velyvis, A. Velyviene, B. Hu, T. Haas, E. Plow, and J. Qin. 2002. A structural mechanism of integrin  $\alpha\text{IIb}\beta 3$  "inside-out" activation as regulated by its cytoplasmic face. *Cell.* 110:587–597.
- Wegener, K.L., A.W. Partridge, J. Han, A.R. Pickford, R.C. Liddington, M.H. Ginsberg, and I.D. Campbell. 2007. Structural basis of integrin activation by talin. *Cell.* 128:171–182.
- Xi, X., R.J. Bodnar, Z. Li, S.C.-T. Lam, and X. Du. 2003. Critical roles for the COOH-terminal NITY and RGT sequences of the integrin  $\beta 3$  cytoplasmic domain in inside-out and outside-in signaling. *J. Cell Biol.* 162:329–339.
- Xi, X., P. Flevaris, A. Stojanovic, A. Chishti, D.R. Phillips, S.C. Lam, and X. Du. 2006. Tyrosine phosphorylation of the integrin  $\beta 3$  subunit regulates beta(3) cleavage by calpain. *J. Biol. Chem.* 281:29426–29430.
- Zou, Z., H. Chen, A.A. Schmaier, R.O. Hynes, and M.L. Kahn. 2007. Structure-function analysis reveals discrete beta3 integrin inside-out and outside-in signaling pathways in platelets. *Blood.* 109:3284–3290.