

# DEMONSTRATION OF MITOCHONDRIAL ATPase ACTIVITY IN FORMALIN-FIXED COLONIC EPITHELIAL CELLS

LUIS R. OTERO-VILARDEBÓ, NATHAN LANE, and GABRIEL C. GODMAN. From the Department of Microbiology, Columbia University, New York

## INTRODUCTION

The intracellular distribution of adenosine triphosphatase (ATPase) activity has been extensively studied by means of two histochemical methods in which the inorganic phosphate (Pi) produced upon the enzymatic hydrolysis of adenosine triphosphate (ATP) is trapped as an insoluble salt (26, 40). The application of these methods in light microscope studies on a variety of unfixed tissues revealed the ATPase activity on the cell membranes, microvilli, and mitochondria (13, 15, 19, 23, 24, 27, 39, 41).

The mitochondrial activity was in no case demonstrable in tissue which had been chemically fixed for more than 90 minutes, and usually for shorter periods (13, 18, 23, 24, 35, 39, 40, 41). Fixation of cell fractions from liver homogenates reduced the mitochondrial ATPase activity by 80 to 90 per cent of the original activity (24). Even very brief fixation in OsO<sub>4</sub> completely inhibited the mitochondrial activity, while not all structural details were adequately preserved for electron microscopy (9). Shortly thereafter the preservation of ultrastructure and enzymatic activity in general was greatly improved by fixation in formalin preceding incubation (16, 38), and subsequently a large number of other aldehydes have been employed for enzyme histochemistry (32). However, prolonged fixation in the aldehydes also inhibits the mitochondrial ATPase. Thus the electron microscope studies on tissues subjected to prolonged chemical fixation have not revealed significant deposits within the mitochondria after incubation for ATPase activity (1, 8, 9, 16, 19, 21, 22, 29, 32, 33, 37, 38).

A reaction product within mitochondria has been observed after incubation of fresh frozen sections (1, 19, 33) and of sections very briefly fixed in formalin (1, 19, 39) or osmium tetroxide (1, 29). In these studies, some ultrastructural details of the tissues have been compromised in order to retain maximal enzymatic activity.

In this note we report some unusual observa-

tions on the mitochondria of rat colonic mucosa incubated to reveal ATPase activity after overnight fixation in 4 per cent formaldehyde.

## MATERIALS AND METHODS

The descending colon from stock male Sherman rats was opened, cut into pieces about 5 × 5 mm, and fixed overnight (approximately 16 hours) at 4°C in 4 per cent formaldehyde in 0.05 M cacodylate buffer and 1 per cent CaCl<sub>2</sub> at pH 7.4 to 7.6. (The fixative had been prepared 4 months before this study was carried out, but the pH was checked immediately before use.) Ten per cent sucrose was added to the fixative for some specimens. Fixation was followed by rinsing in 0.05 M cacodylate buffer with 10 per cent sucrose at pH 7.4 to 7.6 at 4°C for at least six changes (2 hours total). Some tissues were used immediately, but others were kept in the buffer for 1 month at 4°C. Since the ATPase of kidney has been extensively studied, pieces of this tissue were fixed in the same manner for use as controls.

Tissues were then frozen in a mixture of 95 per cent alcohol and dry ice at -70°C, and frozen sections were cut in a cryostat at 30 to 40 μ and collected in buffered sucrose plus 0.05 M MgCl<sub>2</sub> at 4°C. The sections were then transferred into the incubation mixture modified from that of Wachstein and Meisel (40) to the following composition:

	Test	Control	Final conc.
ATP disodium salt (Pabst) 125 mg/100 ml	20 ml	—	8.3 × 10 <sup>-4</sup> M
Tris-Maleate <sup>1</sup> buffer 0.20 M, pH 7.3	20	20	0.08 M
Pb (NO <sub>3</sub> ) <sub>2</sub> 2 %	2	2	2.5 × 10 <sup>-3</sup> M
MgCl <sub>2</sub> , 0.5 M	5	5	0.05 M
Distilled water	3	23	—

<sup>1</sup> The importance of the purity of maleic acid has been recently discussed (3). In these experiments, we have used No. 2927 from Mathieson, Coleman and Bell as recommended by Barka (personal communication).

The components were added in the order listed. The lead solution was added dropwise while the medium was constantly stirred. The final solution was clear or only very slightly cloudy; in either case, it was filtered. Media with any considerable cloudiness or precipitate yielded negative results. The final pH was adjusted to 7.3.

The sections were incubated at 37°C for various lengths of time (5 to 90 minutes). The incubation was followed by rinses in distilled water, very briefly

initially examined unstained, but for most of the studies the sections were stained for 30 to 60 minutes in a saturated solution of uranyl acetate in 50 per cent ethanol. Preparations were examined in an RCA EMU 2E.

## RESULTS AND DISCUSSION

Lead deposits can be seen on the brush border and the lateral membranes of the rat colonic mucosa




FIGURE 1 Rat colonic mucosa fixed overnight in buffered formaldehyde at 4°C and incubated for 40 minutes in ATP-containing medium. The brush border (*BB*) and multivesicular bodies (*MV*) are covered with very heavy deposits. Interdigitations between lateral membranes are greatly accentuated by the lead deposits. All mitochondria exhibit a banded array of lead deposits.  $\times 10,000$ , unstained.

(ca 20 seconds) in 0.005 M acetic acid, and then two additional rinses in distilled water.

Control tissues were incubated for 90 minutes in substrate-free medium or in complete medium containing  $10^{-3}$  M *N* ethyl maleimide.

Tissues to be examined in the electron microscope were postfixed in 2 per cent  $\text{OsO}_4$  buffered with 0.1 M phosphate, pH 7.6, for 45 to 60 minutes at 4°C. The tissues were embedded in Epon after dehydration through graded alcohols and propylene oxide. Sections were cut on a Huxley-Cambridge ultramicrotome and mounted in the usual way. All tissues were

(Fig. 1), as reported by previous workers for a variety of other tissues (1, 9, 16, 18). The interdigitations between adjacent epithelial cells (10, 12) are very well preserved and greatly accentuated by the lead deposits (Figs. 1 and 2).

The very heavy deposits observed over the multivesicular bodies in both the principal and goblet cells (Figs. 1 and 2) may be due to the action of a non-specific hydrolase. Similar deposits have been observed in these multivesicular bodies

after incubation in the presence of  $\beta$ -glycerophosphate at pH 5.0 (2, 25).

The lead deposits encountered within the mitochondria give them a banded appearance like the array of the *cristae mitochondriales* (Figs. 1 to 4). The mitochondrial membranes are not clearly discernible in these preparations, even after prolonged "staining" with lead or uranyl salts. This difficulty in revealing membranes after fixation

in the mitochondrial reaction was observed, but the banded pattern was still easily recognizable (Fig. 4).

The mitochondrial deposits were not present in tissues incubated in substrate-free medium or in complete medium plus  $10^{-3}$  M *N*-ethyl-maleimide (NEM). Nor were they observed when thiamine pyrophosphate (TPP), inosine diphosphate (IDP), or  $\beta$ -glycerophosphate were used as




FIGURE 2 Rat colonic mucosa prepared as described in Fig. 1. The pattern of the intramitochondrial deposits suggest that they cover the *cristae mitochondriales*. Multivesicular bodies (MV) and cell membranes also exhibit reaction product.  $\times 31,000$ . Stained with uranyl acetate for 20 minutes.

with aldehydes has been experienced by other investigators (22, 32). The deposits are apparently confined to the inner mitochondrial membrane and its convolutions, and are lodged upon the cristae (Fig. 3).

ATPase activity of the colonic epithelial mitochondria, but of different intensity, was discerned in three different experiments using the same methods.

When sections cut from pieces of aldehyde-fixed colon, which had been kept for 1 month in buffer at  $4^{\circ}\text{C}$ , were incubated in the medium, a reduction

of the substrate (25). This suggests that the reaction has some specificity.

It might be argued that deposits such as these may arise as the result of the diffusion of the product from a highly reactive site outside the mitochondria. That this is not the case here is suggested by the three following observations. (a) The multivesicular bodies are highly reactive towards  $\beta$ -glycerophosphate, but the mitochondria in their immediate vicinity never show any deposits (2, 25). (b) Similarly, the endoplasmic reticulum of the colonic mucosa which exhibits




FIGURE 3 High magnification of a preparation similar to those in Figs. 1 and 2. The membranes of the mitochondrion are not clearly discernible. Some deposits (arrows) present a scalloped outline. Micrograph was taken by Dr. L. D. Peachey in an RCA EMU<sub>3</sub>F.  $\times 92,000$ . Stained with uranyl acetate for 60 minutes. Insert,  $\times 138,000$ .

high enzymatic activity towards IDP is often seen in very intimate contact with mitochondria, which are always free of deposits (25). (c) The so called  $\beta$  cytomembranes on the basal portion of the cells in the distal tubule of the kidney are known to delimit compartments containing mitochondria and to exhibit ATPase activity (21, 35), but even after prolonged incubation (90 minutes) in this medium no deposits were apparent in the mitochondria (Fig. 5).

The difference in response between the mitochondria in the kidney and colon is not completely understood. It may be a real difference, or result from different sensitivities of the mitochondrial enzymes to formaldehyde fixation.

The nature and significance of mitochondrial ATPase activity is still debated. This activity is usually attributed to the action of two different enzymes, one of which is enhanced by dinitrophenol (DNP) and another which is stimulated by  $Mg^{++}$  ions. These two enzymes have different pH optima and also exhibit different sensitivities toward inhibitors (4, 5, 14, 34). In most of these studies the activity has been found associated with submitochondrial "particles" obtained by the treatment of mitochondria with digitonin (14), deoxycholate (34), or ultrasonication (7). Electron microscopic examination of the "deoxycholate particles" revealed that these consist of membranous structures (42). From this evidence the mitochondrial ATPase is believed to be membrane bound, presumably to the cristae.

More recently a soluble fraction of the ruptured mitochondria has been reported to exhibit properties which may explain the effects of DNP and  $Mg^{++}$  as the action upon a single enzyme (30). This soluble enzyme may represent a further fractionation of the particles described above. Extensive reviews on this subject have been recently published (20, 31).

The methods for histochemical localization of phosphatases in the electron microscope allow only the detection of the site of deposition of the trapped reaction product, *i.e.* lead phosphate. This may or may not coincide with the exact site of the enzyme. The importance of this limitation is heightened by the results which are being currently obtained with the "negative staining" techniques as applied to viruses (6, 28) and mitochondria (11, 28, 36), in which a macromolecular aggregate is embedded in an electron-opaque substance which fills all the space available except that




FIGURE 4 Rat colonic mucosa fixed overnight in buffered formaldehyde, kept in buffered sucrose for 1 month at 4°C, and incubated 40 minutes for ATPase activity. The intramitochondrial deposits are less pronounced than in Figs. 1 to 3, but the banded pattern is still recognizable.  $\times 31,000$ . Unstained.

FIGURE 5 Basal region of cells lining the distal tubule of rat kidney. Tissue fixed overnight in buffered formaldehyde and incubated for 90 minutes for ATPase activity. The so called  $\beta$  cytomembranes exhibit heavy clumped deposits, but the mitochondria (M) in their immediate vicinity are free of deposits.  $\times 18,000$ . Unstained.

occupied by the aggregate itself. Thus if an active enzyme is part of such a macromolecular aggregate, it may preclude any extensive deposition of the reaction product precisely on the enzyme site, and, consequently, the deposits may produce the effect of a negative stain at high levels of resolution.

In summary, an extensive deposition of reaction product has been observed within the mitochondria in apparent relation to the inner membranes after incubation of formaldehyde-fixed colonic mucosa using ATP as a substrate. Similar results were obtained with these tissues after 1-month storage in buffer at 4°C. The mitochondria of formalin-fixed kidney gave no reaction. The colonic mitochondria showed no deposits in the presence of  $\beta$ -glycerophosphate, IDP, TPP, or ATP plus NEM.

The authors are grateful to Dr. L. D. Peachey for his comments, and to Mr. Bill M. Boland for his assistance.

This study was supported by Grant A-817 from the United States Public Health Service.

Mr. Otero-Vilardebó is a Predoctoral Fellow of the United States Public Health Service in the Department of Zoology, Columbia University. He is on leave from the University of Puerto Rico.

Received for publication, June 7, 1963.

#### REFERENCES

1. ASHWORTH, C. T., LUIBEL, F. J., and STEWART, S. C., *J. Cell Biol.*, 1963, 17, 1.
2. BARKA, T., *J. Histochem. and Cytochem.*, 1963, in press.

3. BARKA, T., and ANDERSON, P. J., *J. Histochem. and Cytochem.*, 1962, **10**, 741.
4. BEYER, R. E., *Biochim. et Biophysica Acta*, 1962, **65**, 289.
5. BOS, C. J., and EMMELOT, P., *Biochim. et Biophysica Acta*, 1962, **64**, 21.
6. BRENNER, S., and HORNE, R. W., *Biochim. et Biophysica Acta*, 1960, **34**, 103.
7. COOPER, C., *Biochim. et Biophysica Acta*, 1958, **30**, 484.
8. DEBEYER, J. M., DEMAN, J. H. C., and PERSIJN, J. P., *J. Cell Biol.*, 1962, **13**, 452.
9. ESSNER, E., NOVIKOFF, A. B., and MASSEK, B., *J. Biophysic. and Biochem. Cytol.*, 1958, **4**, 713.
10. FAWCETT, D. W., in *Frontiers in Cytology* (S. L. Palay, editor), 1958, New Haven, Yale University Press.
11. FERNÁNDEZ-MORÁN, H., *Circulation*, 1962, **26**, 1039.
12. FLOREY, H., *Quart. J. Exp. Physiol.*, 1960, **45**, 329.
13. FREIMAN, D. G., and KAPLAN, N., *J. Histochem. and Cytochem.*, 1960, **8**, 159.
14. GAMBLE, J. L., and LEHNINGER, A. L., *J. Biol. Chem.*, 1956, **223**, 921.
15. HORI, S. H., and CHANG, J. P., *J. Histochem. and Cytochem.*, 1963, **11**, 71.
16. KAPLAN, S. E., and NOVIKOFF, A. B., *J. Histochem. and Cytochem.*, 1959, **7**, 295.
17. KIELLY, W. W., and KIELLY, R. K., *J. Biol. Chem.*, 1953, **200**, 213.
18. KRUS, S., ANDRADE, Z. A., and BARKA, T., *J. Histochem. and Cytochem.*, 1961, **9**, 487.
19. LAZARUS, S. S., and BARDEN, H., *J. Histochem. and Cytochem.*, 1962, **10**, 285.
20. LEHNINGER, A. L., and WADKINS, C. L., *Ann. Rev. Biochem.*, 1962, **31**, 47.
21. NOVIKOFF, A. B., *Circulation*, 1962, **26**, 1126.
22. NOVIKOFF, A. B., DETHE, G., BEARD, D., and BEARD, J. W., *J. Cell Biol.*, 1962, **15**, 451.
23. NOVIKOFF, A. B., DRUCKER, J., SHIN, W. Y., and GOLDFISHER, S., *J. Histochem. and Cytochem.*, 1961, **9**, 434.
24. NOVIKOFF, A. B., HAUSMAN, D. H., and POBDER, E., *J. Histochem. and Cytochem.*, 1958, **6**, 61.
25. OTERO-VILARDEBÓ, L. R., LANE, N., and GODMAN, G. C., *J. Histochem. and Cytochem.*, 1963, in press.
26. PADYKULA, H. A., and HERMAN, E., *J. Histochem. and Cytochem.*, 1955, **3**, 161.
27. PADYKULA, H. A., and HERMAN, E., *J. Histochem. and Cytochem.*, 1955, **3**, 170.
28. PARSONS, D. F., *J. Cell Biol.*, 1963, **16**, 620.
29. PERSIJN, J. P., DEAMS, W. T., DEMAN, J. C. H., and MEIJER, A. E. *Histochemie*, 1961, **2**, 372.
30. PULLMAN, M. E., PENEFSKY, H. S., DATTA, A., and RACKER, E., *J. Biol. Chem.*, 1960, **235**, 3322.
31. RACKER, E., *Adv. Enzymol.*, 1961, **23**, 323.
32. SABATINI, D. D., BENSCH, K., and BARNETT, R. J., *J. Cell Biol.*, 1963, **17**, 19.
33. SCARPELLI, D. G., and CRAIG, H. L., *J. Cell Biol.*, 1963, **17**, 279.
34. SIEKEVITZ, P., LOW, H., ERNSTER, L., and LINDBERG, O., *Biochim. et Biophysica Acta*, 1958, **29**, 378.
35. SPATER, H. W., NOVIKOFF, A. B., and MASSECK, B., *J. Biophysic. and Biochem. Cytol.*, 1958, **4**, 765.
36. STOECKENIUS, W., *J. Cell Biol.*, 1963, **17**, 443.
37. TICE, L., and BARNETT, R. J., *J. Cell Biol.*, 1962, **15**, 401.
38. WACHTEL, A., LEHRER, G. M., MAUTNER, W., DAVIS, B. J., and ORNSTEIN, L., *J. Histochem. and Cytochem.*, 1959, **7**, 291.
39. WACHSTEIN, M., BRADSHAW, M., and ORTIZ, J. M., *J. Histochem. and Cytochem.*, 1962, **10**, 65.
40. WACHSTEIN, M., and MEISEL, E., *Am. J. Clin. Path.*, 1957, **27**, 13.
41. WACHSTEIN, M., MEISEL, E., and NIEDZWIEDZ, A., *J. Histochem. and Cytochem.*, 1960, **8**, 387.
42. WATSON, M. L., and SIEKEVITZ, P., *J. Biophysic. and Biochem. Cytol.*, 1956, **2**, 639.